



UMDEPP-014-024

Adinkras, 0-branes, Holoraumy and the SUSY QFT/QM Correspondence

Mathew Calkins^{1†}, D. E. A. Gates^{2†}, S. James Gates, Jr.^{3†}, and Kory Stiffler^{4*}

[†]*Center for String and Particle Theory
Department of Physics, University of Maryland
College Park, MD 20742-4111 USA*

and

^{*}*Department of Chemistry, Physics, and Astronomy
Indiana University Northwest
Gary, Indiana 46408 USA*

ABSTRACT

We propose the recently defined “Holoraumy Tensor” to play a critical role in defining a metric to establish a correspondence between 4D, $\mathcal{N}$ -extended 0-brane-based valise supermultiplet representations and, correspondingly via “SUSY Holography,” on the space of 1D, N -extended network-based adinkras. Using an analogy with the $\mathfrak{su}(3)$ algebra, it is argued the 0-brane holoraumy tensor plays the role of the “d-coefficients” and provides a newly established tool for investigating supersymmetric representation theory.

PACS: 11.30.Pb, 12.60.Jv

Keywords: quantum mechanics, supersymmetry, off-shell supermultiplets

¹mathewpcalkins@gmail.com

²deagates@terpmail.umd.edu

³gatess@wam.umd.edu

⁴kmstiffl@iun.edu

1 Introduction

There are two logical ways to construct one dimensional theories that possess the property of supersymmetry (SUSY) by starting from:

- (a.) a higher dimensional supersymmetrical field theory and applying torus compactification (in the simplest case), or
- (b.) a one dimensional framework and building supersymmetry up without any assumptions.

These two modes of construction will obviously lead to some area of overlap and in this realm some interesting questions and possibilities occur. The former approach obviously will contain remnants of information about the symmetries of the higher dimensional spacetime. How are these data “downloaded” onto the theories constructed from a purely adinkra network-based [1, 2, 3, 4, 5, 6, 7, 8, 9, 10] approach?

On the other hand, we have argued the off-shell or auxiliary field problem for one dimensional linear representation SUSY theories has a solution [11, 12, 13, 14, 15, 16, 17] based on an algebraic structure we have denoted by the symbol $\mathcal{GR}(d, N)$ or “Garden Algebras.” How can this information be “uploaded,” via the area of overlap between the two approaches, in order to more completely analyze the off-shell or auxiliary field problem for higher dimensional SUSY theories?

This area of mutual overlap also suggests the possibility that kinematical information about the higher dimensional SUSY field theories can be encoded into one dimensional adinkra network-based models. We have long called this idea “SUSY Holography.” In a recent paper, [18] there was introduced a new tensor on the space of adinkras that can be defined for all linear representations. We have argued a “holoraumy tensor” (in a consciously mixed provenance of Greek and German language contributions) is the “Rosetta Stone” for the translation of information between one dimensional adinkra network-based models and higher dimensional field theory ones.

Adinkras are graphical networks for 1D, N -extended SUSY representations, and as such at first there would appear no way for them to carry information about spin of higher dimensional theories. In fact, the mechanism known as “spin from isospin” [19] has long been known in the physics literature and a variant on this seem workable for overcoming this objection. Such a mechanism seems to be at work within SUSY holography and this variation of the idea is critical.

Among the adinkra graphs, there is a special class defined by its members possessing ‘height’ assignments of nodes solely to two distinct levels. Members of this

restricted class of adinkras have been given the name of “valise adinkras.” It had long been the suspicion of one of the authors (SJG) that valise adinkras are in some sense more elementary than all others. In recent works, the sense in which the valise adinkras are more elementary has come sharply into focus. When the spinorial derivatives of one dimensional SUSY theories are evaluated on the fields of a valise adinkra in a specified way this leads to the appearance a tensor (that shares simultaneous attributes of the usual Riemann curvature tensor and the usual torsion tensor tensor). This new tensor is the “holoraumy” tensor.

Since the concept of the “holoraumy” tensor in supersymmetric representation theory is a new one, to make it more familiar it is useful to review an analogy from group theory. The Lie algebra $\mathfrak{su}(3)$ provides a way to do this. We use the Gell-Mann matrix representation of the $\mathfrak{su}(3)$ generators and denote these by $\mathbf{t}_A$ leading to the familiar structure constants $f_{AB}{}^C$

$$[\mathbf{t}_A, \mathbf{t}_B] = i f_{AB}{}^C \mathbf{t}_C, \quad (1)$$

of the algebra. Next we define eight new matrices via the definitions $\tilde{\mathbf{t}}_A = -\mathbf{t}_A^*$ and find these new matrices satisfy

$$[\tilde{\mathbf{t}}_A, \tilde{\mathbf{t}}_B] = i f_{AB}{}^C \tilde{\mathbf{t}}_C, \quad (2)$$

and as this is identical to (1) these eight new matrices also form a representation of the generators of $\mathfrak{su}(3)$. We can introduce a more compact notation by further defining $\mathbf{t}_A^{(\#1)} \equiv \mathbf{t}_A$ and $\mathbf{t}_A^{(\#2)} \equiv \tilde{\mathbf{t}}_A$, so quadratic Casimir operators for each representation are define by

$$C_{(2)}(\mathcal{R}) = \sum_{A=1}^8 \mathbf{t}_A^{(\mathcal{R})} \mathbf{t}_A^{(\mathcal{R})}, \quad (3)$$

which on both takes the value of $16/3$ times the 3×3 identity matrix. Yet, these two representations of the generators of the $\mathfrak{su}(3)$ algebra are not equivalent to one another. That is there does not exist a unitary transformation that relates one to the other. One way to tease out this difference is by starting with the introduction “d-coefficients” for each representation

$$\{\mathbf{t}_A^{(\mathcal{R})}, \mathbf{t}_B^{(\mathcal{R})}\} = \frac{1}{3} \delta_{AB} \mathbf{I}_{3 \times 3} + d_{AB}^{(\mathcal{R})C} \mathbf{t}_C^{(\mathcal{R})}, \quad (4)$$

with $d_{AB}^{(\#1)C} \equiv d_{AB}{}^C$ and $d_{AB}^{(\#2)C} \equiv \tilde{d}_{AB}{}^C$. The index $\mathcal{R}$ plays the role of a “representation label.” We are free to define a metric (not the Killing metric) $\tilde{g}(\mathcal{R}, \mathcal{R}')$ on this representation space via the equation

$$\tilde{g}(\mathcal{R}, \mathcal{R}') = \mathcal{N}_0 \sum_{A,B,C} d_{AB}^{(\mathcal{R})C} d_{AB}^{(\mathcal{R}')C} \quad (5)$$

where $\mathcal{N}_0$ is a normalization constant. The form of this metric leads to a 2×2 matrix

$$\tilde{g}(\mathcal{R}, \mathcal{R}') = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, \quad (6)$$

after using (1) - (5) and an appropriate choice of $\mathcal{N}_0$. Thus the metric defined by (5) is capable of distinguishing between the two representations of $\text{su}(3)$. To construct this new metric required the d-coefficients and we assert holonomy tensors are the analogs for supersymmetric representations.

Following the first lines of the introduction above about the two ways to construct 1D, N -extended supermultiplets, there exist two logically distinct constructions of the holonomy tensor. One follows from the line of argument utilizing higher dimensional theory as a starting point and the other from the solely one dimensional vantage. On equating these two different definitions (a condition we have called the Adinkra/ γ -matrix Equation previously [24, 25]) one defines a projection operator $\mathcal{P}$ capable of mapping higher dimensional 0-brane-based holonomy results into one dimensional network-based holonomy results. In the following we will explore additional properties of these holonomy tensors in some examples.

2 Network-Based Valise Adinkra Representations

We have established a well-defined methodology [3, 4, 5] for creating a set of graphs (called ‘adinkras’) capable of providing a basis for constructing linear representations of 1D, N -extended off-shell supermultiplets. This methodology is *totally* divorced from any higher dimensional concepts, and accordingly the method contains no a priori relations to nor assumptions about symmetries in higher dimensions. Conceptually this approach can be based on networks and their associated “adinkra” adjacency matrices. Three examples of these are shown below.




Figure # 1: Three adinkra graphs

Of these three adinkras, only the final one has the valise property - all open nodes appear at the same height in the diagram and all closed nodes appear at the same height also but which is distinct from the height of the open nodes. This third adinkra is a member of the subclass of ‘valise adinkras.’ Though the first two diagrams are not valise adinkras, through a series of operations designed for the lowering of nodes successively, these two adinkras can also be brought to the form of valise adinkras. Thus, all three adinkras shown in the Fig. # 2 are valise adinkras.




Figure # 2: Three valise adinkra graphs

The parameter N describes the number of equivalence classes of the links. In the adinkras, it is shown by the N distinct colors used to draw the diagram. Any two links in the same equivalence class possess the same color.

There is a subtlety we should mention before moving on. The definition of the adinkra graphs given above is a coordinate-independent one. In order to make any calculation with an adinkra, one must choose a set of coordinate basis elements. This has been done for the three adinkras $\mathcal{R} = \#1, \#2$, and $\#3$ shown in Fig. # 3




Figure # 3: Three valise adinkra graphs with node assignment

where the values of $\mathcal{R}$ can be used to describe which adinkra is under consideration.

One of these choices of bases is the reference basis to which the open nodes are defined. A second choice of these bases is the reference basis to which the closed nodes are defined. The final choice of basis is the one to which the links are referred. The first basis choice is a choice of partitioning of the numbers $1 \dots d$ among the bosonic nodes. The second basis choice is a choice of partitioning of the numbers $1 \dots d$ among the fermionic nodes. The final choice one of the numbers $1 \dots N$ among the classes of the links.

Each open node represents a bosonic function (or any of its derivatives) of τ (a time-like parameter) and collectively we denote these by Φ_i with $i = 1, \dots, d$ and similarly, each closed node represents a fermionic function (or its derivatives) of τ and collectively we denote these by $\Psi_{\hat{k}}$ with $\hat{k} = 1, \dots, d$. As the construction of network-based adinkra graphs begins with the partition of an appropriate hypercube, the equality in the number of bosons and fermions follows as a consequence. As in our example above, any adinkra which is not in the form of a valise can be brought to that form by judicious lowering of nodes.

Valise adinkras are special in the form of the supercovariant derivative takes when acting on the bosons and fermions in the supermultiplet can be expressed as

$$D_I \Phi_i = i (L_I)_{i\hat{k}} \Psi_{\hat{k}} \quad , \quad D_I \Psi_{\hat{k}} = (R_I)_{\hat{k}i} \partial_\tau \Phi_i \quad , \quad (1)$$

in terms of constants $(L_I)_{i\hat{k}}$ and $(R_I)_{\hat{k}i}$ where $I = 1, \dots, N$. As we have emphasized many times before (see for example [11, 12, 13, 14, 15, 16, 17]), when an appropriate set of algebraic conditions are imposed on $(L_I)_{i\hat{k}}$ and $(R_I)_{\hat{k}i}$, the equations in (1) imply

$$\{D_I, D_J\} \Phi_i = i 2 \delta_{IJ} \partial_\tau \Phi_i \quad , \quad \{D_I, D_J\} \Psi_{\hat{k}} = i 2 \delta_{IJ} \partial_\tau \Psi_{\hat{k}} \quad , \quad (2)$$

i.e. the collections of functions $(\Phi_i, \Psi_{\hat{k}})$ form a supermultiplet.

Using the basis described by the adinkras in Fig. # 3, we find in the respective cases of the three adinkras

	$\mathcal{R} = \# 1$	$\mathcal{R} = \# 2$	$\mathcal{R} = \# 3$
<i>RED</i>	$(10)_b (243)$	$(10)_b (1243)$	$(14)_b (234)$
<i>BLACK</i>	$(0)_b (142)$	$(6)_b (1342)$	$(2)_b (143)$
<i>GREEN</i>	$(12)_b (123)$	$(12)_b (23)$	$(4)_b (124)$
<i>BLUE</i>	$(6)_b (134)$	$(0)_b (14)$	$(8)_b (132)$

Table # 1: Adinkra Link Color & Cycles in L-matrices

where we have used a shorthand notation introduced in [20, 21] to denote 4×4 matrices. The final step needed to determine the relations of the L-matrices to the

graph is the assignment of a ‘rainbow’ (as described in [22]) where each color is assigned to a number. In the following we make the choices *red* = 1, *black* = 2, *green* = 3, and *blue* = 4.

3 Network-Based Fermionic Holoraumy Tenors

When an adinkra is a valise, there is another set of equations that are satisfied. These take the forms

$$\begin{aligned} [D_I, D_J] \Phi_i^{(\mathcal{R})} &= -i 2 [\mathcal{B}_{IJ}^{(\mathcal{R})}]_{ik} \partial_\tau \Phi_k^{(\mathcal{R})} , \\ [D_I, D_J] \Psi_{\hat{k}}^{(\mathcal{R})} &= -i 2 [\mathcal{F}_{IJ}^{(\mathcal{R})}]_{\hat{k}\hat{\ell}} \partial_\tau \Psi_{\hat{\ell}}^{(\mathcal{R})} . \end{aligned} \quad (3)$$

In the work of [18], we have given the quantities $[\mathcal{B}_{IJ}^{(\mathcal{R})}]_{ik}$ and $[\mathcal{F}_{IJ}^{(\mathcal{R})}]_{\hat{k}\hat{\ell}}$ respectively the names of the “bosonic holoraumy tensor” and the “fermionic holoraumy tensor.” We include the “representation label” $\mathcal{R}$ within these equations to recognize that each of the three adinkras in Fig. # 3 has distinct L-matrices (and R-matrices) depending on the adinkra under consideration.

From (1) and (3), it follows

$$\begin{aligned} [D_I, D_J] \Phi_i^{(\mathcal{R})} &= -i 2 [\mathcal{B}_{IJ}^{(\mathcal{R})}]_{ik} \partial_\tau \Phi_k^{(\mathcal{R})} = -i 2 [V_{IJ}^{(\mathcal{R})}]_i{}^k \left(\partial_\tau \Phi_k^{(\mathcal{R})} \right) , \\ [D_I, D_J] \Psi_{\hat{k}}^{(\mathcal{R})} &= -i 2 [\mathcal{F}_{IJ}^{(\mathcal{R})}]_{\hat{k}\hat{\ell}} \partial_\tau \Psi_{\hat{\ell}}^{(\mathcal{R})} = -i 2 [\tilde{V}_{IJ}^{(\mathcal{R})}]_{\hat{k}}{}^{\hat{\ell}} \left(\partial_\tau \Psi_{\hat{\ell}}^{(\mathcal{R})} \right) , \end{aligned} \quad (4)$$

where

$$\begin{aligned} [V_{IJ}^{(\mathcal{R})}]_i{}^k &= \frac{1}{2} \left[(L_I^{(\mathcal{R})})_i{}^j (R_J^{(\mathcal{R})})_j{}^k - (L_J^{(\mathcal{R})})_i{}^j (R_I^{(\mathcal{R})})_j{}^k \right] , \\ [\tilde{V}_{IJ}^{(\mathcal{R})}]_{\hat{k}}{}^{\hat{\ell}} &= \frac{1}{2} \left[(R_I^{(\mathcal{R})})_{\hat{k}}{}^j (L_J^{(\mathcal{R})})_j{}^{\hat{\ell}} - (R_J^{(\mathcal{R})})_{\hat{k}}{}^j (L_I^{(\mathcal{R})})_j{}^{\hat{\ell}} \right] . \end{aligned} \quad (5)$$

On the space of the three different adinkra representation seen in Fig. # 3, we can define a ‘metric’ via the definition

$$\mathcal{G}[(\mathcal{R}), (\mathcal{R}')] = \frac{1}{48} \sum_{I,J} \text{Tr} \left[\tilde{V}_{IJ}^{(\mathcal{R})} \tilde{V}_{IJ}^{(\mathcal{R}')} \right] , \quad (6)$$

so a few computations reveal a 3×3 matrix

$$\mathcal{G}[(\mathcal{R}), (\mathcal{R}')] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -\frac{1}{3} \\ 0 & -\frac{1}{3} & 1 \end{bmatrix} . \quad (7)$$

In the work of [20] we showed there exists 1,536 ways to construct adinkra networks of four colors, four open nodes, and four closed nodes. The ones shown in Fig. # 3

are only three of this multitude. So in principle, for each of these 1,536 networks one could assign a value of $\mathcal{R}$ so $\# 1 \leq \mathcal{R} \leq \# 1,536$ and for each of these values there exists a calculable $\tilde{V}$ -matrix. Thus, the metric defined by (6) can be extended over the entirety of a 1,536 dimensional space.

Let us close by noting in the discussion above, there is no reference at all to higher dimensional supersymmetrical theories nor their symmetries.

4 0-Brane-Based Valise Supermultiplets

By now in a number of our past works (e. g. [23]), we have discussed the process by which we pass from a standard Lorentz-covariant four dimensional formulation of a supermultiplet to obtain a corresponding QM supermultiplet in valise form, where equations with the following form are valid

$$D_a \Phi_\Lambda = i (L_\Lambda)_a^{\hat{\Lambda}} \Psi_{\hat{\Lambda}} \quad , \quad D_a \Psi_{\hat{\Lambda}} = (R^\Lambda)_{\hat{\Lambda} a} \partial_\tau \Phi_\Lambda \quad , \quad (8)$$

and where the explicit forms of Φ_Λ and $\Psi_{\hat{\Lambda}}$ vary for each multiplet. In particular for the CS we have

$$\Phi_\Lambda = (A, B, F, G) \quad , \quad \Psi_{\hat{\Lambda}} = (\psi_a) \quad , \quad (9)$$

for the VS we have

$$\Phi_\Lambda = (A_m, d) \quad , \quad \Psi_{\hat{\Lambda}} = (\lambda_a) \quad , \quad (10)$$

for the TS we have

$$\Phi_\Lambda = (\varphi, B_{mn}) \quad , \quad \Psi_{\hat{\Lambda}} = (\chi_a) \quad , \quad (11)$$

for the AVS we have

$$\Phi_\Lambda = (U_m, \tilde{d}) \quad , \quad \Psi_{\hat{\Lambda}} = (\tilde{\lambda}_a) \quad , \quad (12)$$

and for the ATS we have

$$\Phi_\Lambda = (\tilde{\varphi}, \tilde{B}_{mn}) \quad , \quad \Psi_{\hat{\Lambda}} = (\tilde{\chi}_a) \quad . \quad (13)$$

These equations demonstrate the indices Λ range over distinct bosonic representations of the $SO(1, 3)$. While in all of these examples the $\hat{\Lambda}$ index corresponds to solely spinor indices of $SO(1, 3)$, for more general supermultiplets (e. g. supergravity, etc.) this

index can range over distinct fermionic representations of $SO(1, 3)$. Below the explicit forms of the L_Λ and R^Λ coefficients can be read out for the cases of

(a.) Chiral Supermultiplet (CS);

$$\begin{aligned} D_a A &= \psi_a \quad , \quad D_a B = i(\gamma^5)_a{}^b \psi_b \quad , \quad D_a F = (\gamma^0)_a{}^b \psi_b \quad , \quad D_a G = i(\gamma^5 \gamma^0)_a{}^b \psi_b \quad , \\ D_a \psi_b &= i(\gamma^0)_{ab} (\partial_\tau A) - (\gamma^5 \gamma^0)_{ab} (\partial_\tau B) - iC_{ab} (\partial_\tau F) + (\gamma^5)_{ab} (\partial_\tau G) \quad , \end{aligned} \quad (14)$$

(b.) Vector Supermultiplet (VS);

$$\begin{aligned} D_a A_m &= (\gamma_m)_a{}^b \lambda_b \quad , \quad D_a d = i(\gamma^5 \gamma^0)_a{}^b \lambda_b \quad , \\ D_a \lambda_b &= -i(\gamma^0 \gamma^m)_{ab} (\partial_\tau A_m) + (\gamma^5)_{ab} (\partial_\tau d) \quad , \end{aligned} \quad (15)$$

(c.) Tensor Supermultiplet (TS);

$$\begin{aligned} D_a \varphi &= \chi_a \quad , \quad D_a B_{mn} = -\frac{1}{4}([\gamma_m, \gamma_n])_a{}^b \chi_b \quad , \\ D_a \chi_b &= i(\gamma^0)_{ab} \partial_\tau \varphi - i\frac{1}{2}(\gamma^0 [\gamma^m, \gamma^n])_{ab} \partial_\tau B_{mn} \quad , \end{aligned} \quad (16)$$

(d.) Axial vector Supermultiplet (AVS); and

$$\begin{aligned} D_a U_m &= i(\gamma^5 \gamma_m)_a{}^b \tilde{\lambda}_b \quad , \quad D_a \tilde{d} = -(\gamma^0)_a{}^b \partial_\tau \tilde{\lambda}_b \quad , \\ D_a \tilde{\lambda}_b &= (\gamma^5 \gamma^0 \gamma^m)_{ab} (\partial_\tau U_m) + iC_{ab} \tilde{d} \quad , \end{aligned} \quad (17)$$

(e.) Axial tensor Supermultiplet (ATS)

$$\begin{aligned} D_a \tilde{\varphi} &= i(\gamma^5)_a{}^b \tilde{\chi}_b \quad , \quad D_a \tilde{B}_{mn} = -i\frac{1}{4}(\gamma^5 [\gamma_m, \gamma_n])_a{}^b \tilde{\chi}_b \quad , \\ D_a \tilde{\chi}_b &= -(\gamma^0 \gamma^5)_{ab} \partial_\tau \tilde{\varphi} + \frac{1}{2}(\gamma^0 \gamma^5 [\gamma^m, \gamma^n])_{ab} \partial_\tau \tilde{B}_{mn} \quad . \end{aligned} \quad (18)$$

From (8) it follows

$$\begin{aligned} [D_a, D_b] \Phi_\Lambda &= -i \left[(L_\Lambda)_a{}^{\hat{\Lambda}} (R^\Delta)_{\hat{\Lambda} b} - (L_\Lambda)_b{}^{\hat{\Lambda}} (R^\Delta)_{\hat{\Lambda} a} \right] \partial_\tau \Phi_\Delta \quad , \\ [D_a, D_b] \Psi_{\hat{\Lambda}} &= -i \left[(R^\Delta)_{\hat{\Lambda} a} (L_\Delta)_b{}^{\hat{\Delta}} - (R^\Delta)_{\hat{\Lambda} b} (L_\Delta)_a{}^{\hat{\Delta}} \right] \partial_\tau \Psi_{\hat{\Delta}} \quad . \end{aligned} \quad (19)$$

The first one of these is a calculation that requires taking the antisymmetric part of some products of strings of γ -matrices. The second one of these is of the form of Fierz identities of some products of strings of γ -matrices. So the character of the coefficients appearing is very different from the similar looking coefficients in (5). At a minimum as the coefficients in (19) are constructed from γ -matrices, information about the symmetries of spacetime are inherited in the 1D supermultiplets discussed in this chapter. The question is, “How is such information to be recovered in the supermultiplets constructed from network-based adinkras?”

5 0-Brane Fermionic Holoraumy Tenors

Direct calculations show the operator equation

$$\{D_a, D_b\} = i2(\gamma^0)_{ab}\partial_\tau, \quad (20)$$

is satisfied on all fields in (14) - (18) as expected by SUSY. However, as each of the equations in (14) - (18) describes a valise supermultiplet, there is something else we can do. The anti-commutator in (20) can be replaced by a commutator.

Our conventions are such that when using the commutator (not the anti-commutator as in (20)) of two spinor covariant derivatives evaluated on the fermions of a covariant valise supermultiplet we write

$$[D_a, D_b] = -i2(\mathcal{F}_{ab})\partial_\tau, \quad (21)$$

where $(\mathcal{F}_{ab})$ denotes a set of representation dependent constants (these will be discussed in more detail via examples below) and when evaluated on the bosons of a valise supermultiplet we write

$$[D_a, D_b] = -i2(\mathcal{B}_{ab})\partial_\tau, \quad (22)$$

where $(\mathcal{B}_{ab})$ denotes a set of representation dependent constants once more. In the remainder of this chapter, we are going to focus on the fermionic holoraumy tensor in (21) and defer discussion of the bosonic holoraumy tensor in (22) until an appendix.

For the fermionic fields in (14) - (18) we find

$$\begin{aligned} (\mathcal{F}_a^{b(CS)})_c^d &= -(\gamma^5\gamma^m)_a^b(\gamma^5\gamma^0\gamma_m)_c^d, \\ (\mathcal{F}_a^{b(VS)})_c^d &= +\delta_a^b(\gamma^0)_c^d + (\gamma^5)_a^b(\gamma^5\gamma^0)_c^d + (\gamma^5\gamma^0)_a^b(\gamma^5)_c^d, \\ (\mathcal{F}_a^{b(TS)})_c^d &= -\delta_a^b(\gamma^0)_c^d + (\gamma^5)_a^b(\gamma^5\gamma^0)_c^d - (\gamma^5\gamma^0)_a^b(\gamma^5)_c^d, \\ (\mathcal{F}_a^{b(AVS)})_c^d &= -\delta_a^b(\gamma^0)_c^d - (\gamma^5)_a^b(\gamma^5\gamma^0)_c^d + (\gamma^5\gamma^0)_a^b(\gamma^5)_c^d, \\ (\mathcal{F}_a^{b(ATS)})_c^d &= +\delta_a^b(\gamma^0)_c^d - (\gamma^5)_a^b(\gamma^5\gamma^0)_c^d - (\gamma^5\gamma^0)_a^b(\gamma^5)_c^d. \end{aligned} \quad (23)$$

A further set of identities given by

$$\begin{aligned} (\mathcal{F}_a^{b(CS)})_c^d &= (\gamma^5)_c^e(\mathcal{F}_a^{b(CS)})_e^f(\gamma^5)_f^d, \\ (\mathcal{F}_a^{b(AVS)})_c^d &= (\gamma^5)_c^e(\mathcal{F}_a^{b(VS)})_e^f(\gamma^5)_f^d, \\ (\mathcal{F}_a^{b(ATS)})_c^d &= (\gamma^5)_c^e(\mathcal{F}_a^{b(TS)})_e^f(\gamma^5)_f^d, \end{aligned} \quad (24)$$

are also valid. These equations have clear interpretations. Under the conjugation operation defined by a similarity transformation using γ^5 , the holoraumy of the valise

chiral supermultiplet is self-conjugate, while the holoraumy of the vector (tensor) valise supermultiplet is conjugate to the holoraumy of the axial-vector (axial-tensor) valise supermultiplet.

A second type of conjugation of the covariant holoraumy tensors can be defined by replacing the γ^5 in (24) by γ^0 . From the equations above (23), it also follows

$$\begin{aligned}
(\gamma^0)_c^e (\mathcal{F}_a^{b(CS)})_e^f (\gamma^0)_f^d &= + (\gamma^5 \gamma^m)_a^b (\gamma^5 \gamma^0 \gamma_m)_c^d, \\
(\gamma^0)_c^e (\mathcal{F}_a^{b(VS)})_e^f (\gamma^0)_f^d &= - \delta_a^b (\gamma^0)_c^d + (\gamma^5)_a^b (\gamma^5 \gamma^0)_c^d + (\gamma^5 \gamma^0)_a^b (\gamma^5)_c^d, \\
(\gamma^0)_c^e (\mathcal{F}_a^{b(TS)})_e^f (\gamma^0)_f^d &= + \delta_a^b (\gamma^0)_c^d + (\gamma^5)_a^b (\gamma^5 \gamma^0)_c^d - (\gamma^5 \gamma^0)_a^b (\gamma^5)_c^d, \\
(\gamma^0)_c^e (\mathcal{F}_a^{b(AVS)})_e^f (\gamma^0)_f^d &= + \delta_a^b (\gamma^0)_c^d - (\gamma^5)_a^b (\gamma^5 \gamma^0)_c^d + (\gamma^5 \gamma^0)_a^b (\gamma^5)_c^d, \\
(\gamma^0)_c^e (\mathcal{F}_a^{b(ATS)})_e^f (\gamma^0)_f^d &= - \delta_a^b (\gamma^0)_c^d - (\gamma^5)_a^b (\gamma^5 \gamma^0)_c^d - (\gamma^5 \gamma^0)_a^b (\gamma^5)_c^d.
\end{aligned} \tag{25}$$

Given the results in (23) and (25), we can define an inner product on the covariant 0-brane space of representations. With the covariant 0-brane reduced supermultiplets (CS) , (VS) , (TS) , (AVS) , and (ATS) , we can introduce a “representation index” denoted by $(\mathcal{R})$ that simply takes on each of the values (CS) , (VS) , (TS) , (AVS) , and (ATS) . We can regard these as five distinct vectors in a representation space. We can introduce a inner product on this vector space. Let $(\mathcal{R})$ and $(\mathcal{R}')$ denote any two of these representation, then we can introduce a inner product, denoted by $\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')]$ on this space through the definition:

$$\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')] = \tilde{\mathcal{G}}[(\mathcal{R}'), (\mathcal{R})] = \frac{1}{48} (\gamma^0)_c^e (\mathcal{F}_{ab}^{(\mathcal{R})})_e^f (\gamma^0)_f^d (\mathcal{F}^{ab(\mathcal{R}')})_d^c. \tag{26}$$

Several direct calculations lead to a result cast into the form of a 5×5 matrix

$$\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')] = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} \\ 0 & -\frac{1}{3} & 1 & -\frac{1}{3} & -\frac{1}{3} \\ 0 & -\frac{1}{3} & -\frac{1}{3} & 1 & -\frac{1}{3} \\ 0 & -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} & 1 \end{bmatrix}, \tag{27}$$

where the rows and columns are respectively labeled by (CS) , (VS) , (TS) , (AVS) , and (ATS) . Given the definition in (26), we see the (CS) , (VS) , (TS) , (AVS) , and (ATS) representations are equivalent to unit vectors since,

$$\begin{aligned}
\tilde{\mathcal{G}}[(CS), (CS)] &= \tilde{\mathcal{G}}[(VS), (VS)] &= \tilde{\mathcal{G}}[(TS), (TS)] &= \\
&= \tilde{\mathcal{G}}[(AVS), (AVS)] &= \tilde{\mathcal{G}}[(ATS), (ATS)] &= 1.
\end{aligned} \tag{28}$$

It is striking that the only entries in (27) are $(-1/3, 0, 1)$ and these are exactly the values found in the metric introduced in (7) for the three network-based adinkras.

Furthermore, we can define an angle between any two of the representations $(\mathcal{R})$ and $(\mathcal{R}')$ via the definition

$$\cos \{\theta[(\mathcal{R}), (\mathcal{R}')]\} = \frac{\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')]}{\sqrt{\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R})]}\sqrt{\tilde{\mathcal{G}}[(\mathcal{R}'), (\mathcal{R}')]} \quad . \quad (29)$$

We have thus obtained a geometrical viewpoint of the five minimal off-shell 4D, $\mathcal{N} = 1$ supersymmetry representations. The unit vector representing the CS representation is orthogonal to the unit vectors representing the VS, TS, AVS, and ATS representations. The angles between the remaining representations can be read from the matrix given in (27) to have a common value of θ_{TV} where

$$\cos(\theta_{TV}) = -\frac{1}{3} \quad . \quad (30)$$

On the subspace of the CS, VS, and TS representations, this is illustrated as given previously in the work of [24, 25].




Figure # 4: Illustration of the CS-VS-TS subspace using the $\tilde{\mathcal{G}}$ metric

Though the diagram above in Fig. # 4 is the same as the one in [24, 25], the interpretation here is very different. In the previous work, this diagram was based on the calculations using network-based adinkras (see Fig. # 3) and using the $\mathcal{G}$ -metric defined by equation (6). The $\tilde{\mathcal{G}}$ -metric was not known then and the image (Fig. # 4) is calculated based on (26) and 0-brane-based valise supermultiplets. The fact that the two images are identical is due to SUSY holography.

It is possible on more general 4D, $\mathcal{N} = 1$ supermultiplets the fermionic holonomy tensor could be of the form

$$\begin{aligned}
(\mathcal{F}_a^{b(\mathcal{R})})_c^d &= (\gamma^5 \gamma^m)_a^b \tilde{\ell}_{mn} (\gamma^5 \gamma^0 \gamma^n)_c^d \\
&+ \delta_a^b \ell_{11} (\gamma^0)_c^d + \delta_a^b \ell_{12} (\gamma^5 \gamma^0)_c^d + \delta_a^b \ell_{13} (\gamma^5)_c^d \\
&+ (\gamma^5)_a^b \ell_{21} (\gamma^0)_c^d + (\gamma^5)_a^b \ell_{22} (\gamma^5 \gamma^0)_c^d + (\gamma^5)_a^b \ell_{23} (\gamma^5)_c^d \\
&+ (\gamma^5 \gamma^0)_a^b \ell_{31} (\gamma^0)_c^d + (\gamma^5 \gamma^0)_a^b \ell_{32} (\gamma^5 \gamma^0)_c^d + (\gamma^5 \gamma^0)_a^b \ell_{33} (\gamma^5)_c^d \\
&+ \left[\delta_a^b \widehat{\ell}_{1m} + (\gamma^5)_a^b \widehat{\ell}_{2m} + (\gamma^5 \gamma)_a^b \widehat{\ell}_{3m} \right] (\gamma^5 \gamma^0 \gamma^m)_c^d \\
&+ (\gamma^5 \gamma^0 \gamma^m)_a^b \left[\bar{\ell}_{m1} \delta_c^d + \bar{\ell}_{m2} (\gamma^5 \gamma^0)_c^d + \bar{\ell}_{m3} (\gamma^5)_c^d \right] ,
\end{aligned} \tag{31}$$

containing the 36 real parameters ℓ_{mn} , $\tilde{\ell}_{mn}$, $\widehat{\ell}_{mn}$ and $\bar{\ell}_{mn}$. It will require further studies of 0-brane valise supermultiplets to determine if such representations exist.

The results in this chapter provide yet another confirmation of the idea of ‘‘SUSY Holography.’’ It is possible to define other inner products on the space of 0-brane SUSY representations. However, the one in (26) matched with a proposal made previously in the work of [24, 25] and is the same (up to a choice of normalization) at that given in [24, 25]. In (26), the representations are those obtained by 0-brane reduction from higher dimensional supermultiplets. In (6), the representations are those obtained directly from network-based adinkra graphs. The results for the angles between the different representations match perfectly.

6 An Intermezzo For Matchings

By starting from the network viewpoint one has a set of bosons Φ_i corresponding to half the nodes in a network and a set of fermions $\Psi_{\hat{k}}$ corresponding to the other half. The i -indices on the bosons Φ_i (the $\hat{k}$ -indices on the fermions $\Psi_{\hat{k}}$) in the equations

$$D_I \Phi_i = i (L_I)_{i\hat{k}} \Psi_{\hat{k}} \quad , \quad D_I \Psi_{\hat{k}} = (R_I)_{\hat{k}i} \partial_\tau \Phi_i \quad , \tag{32}$$

simply counts the bosons (fermions). As such, they *a priori contain no information about space-time symmetries*.

On the other hand, starting from a field theoretical viewpoint, one has a set of bosons Φ_Λ and fermions $\Psi_{\widehat{\Lambda}}$. The Λ -indices on the bosons Φ_Λ (the $\widehat{\Lambda}$ -indices on the fermions $\Psi_{\widehat{\Lambda}}$) in the equations

$$D_a \Phi_\Lambda = i (L_\Lambda)_a^{\widehat{\Lambda}} \Psi_{\widehat{\Lambda}} \quad , \quad D_a \Psi_{\widehat{\Lambda}} = (R^\Lambda)_{\widehat{\Lambda}a} \partial_\tau \Phi_\Lambda \quad , \tag{33}$$

are spacetime indices (i. e. scalar, vector, etc. for Λ and spinor, etc. for $\widehat{\Lambda}$) and thus provide explicit representations of space-time symmetries.

So though it is true there are many more constructions possible involving the adjacency matrices associated with networks realizations of off-shell 1D, N -extended SUSY, a subset of the models described by (32) must in fact be identical to the models described by (33). For this subset, it follows there must be “hidden” relations between the network-based L_I and R_I matrices on one side of a correspondence and the L_Λ and R^Λ coefficients on the other side. This implies within the subset, the system described by (32) must be a hologram of the system described by (33). Hence, SUSY holography must be realized.

In the work of [24, 25], an equation called the “Adinkra/ γ -matrix Equation” was presented without comment. From our present vantage point, this equation can be explained more fully. As we saw in chapters two and three, beginning with network-based valise adinkras it is easy to calculate their associated V -matrices and $\widetilde{V}$ -matrices.

In chapters four and five, it was shown that application of 0-brane reduction and node-lowering field redefinitions can be applied to 4D, $\mathcal{N} = 1$ supermultiplets and result in equations that have the same general form as those that arise from a valise adinkra. This allows the calculations of the covariant holoraumy tensors $\mathcal{B}_{ab}$ and $\mathcal{F}_{ab}$. The next task must be to unearth the mathematical tools permitting supermultiplets from networks to be mapped onto supermultiplets obtained by 0-brane reduction. Three such tools have been identified for this purpose.

(a.) The Adinkra/ γ -matrix Equation

In the work of [24, 25], the assertion appears there exist a projection operator $\mathcal{P}$ such that

$$\mathcal{P}(\mathcal{F}_{ab}) = [\mathcal{F}_{IJ}] \tag{34}$$

so the components of the covariant fermionic holoraumy tensor $\mathcal{F}_{ab}$ can be set equal to the entries of the $\widetilde{V}$ -matrices. This provides a sort of “Rosetta Stone” that performs the translation between the higher dimensional SUSY field theory on one side of the correspondence and the “Garden Algebra,” network-based adinkras, and codes on the other side.

According to the observations in [22], this translation on the adinkra side will also be important to understand how supermultiplets emerge from Riemann surfaces and Bayli pairs.

This mapping is not one-to-one as was understood from our initial discussions of adinkras. In the work of [1], there appears an image of a pyramid meant to elicit the notion that the number of distinct SUSY representation decreases as the dimension of the superspace increases. This was not meant in a metaphorical sense.

One way to see all of this playing out is in the context of the minimal off-shell representations of 4D, $\mathcal{N} = 1$ supermultiplets. Counting variants [30], there are only eight distinct such supermultiplets. On the other hand, if one does not permit any similarity transformations in counting all possible four color, four-open node and four-closed node adinkras there are 1,536 such objects. So on this subclass of objects the operator $\mathcal{P}$ projects the eight supermultiplets into the ‘sea’ of 1,536 adinkras.

(b.) Coxeter Algebra/Hodge Duality Orbit Matching

In the work of [20], it was shown using the action of the Hodge duality map acting on the fields in equations (33) leads to shadows of these maps that can be detected on the fields in equations (32) and the L-matrices in Table # 1. With this identification, Coxeter algebras were found to play a hidden role in organizing the 1,536 networks so the action of the Hodge $*$ -operator on the 0-brane-based equations of (32) induces the existence of shadow of the Hodge $*$ -operator on the adinkra network-based equations of (33) implying the equation $\mathcal{P}(*\mathcal{F}_{ab}) = [* \mathcal{F}_{IJ}]$.




Figure # 5: Illustration of Hodge $*$ -operator

The diagram in Fig. # 5 shows the orbit matching for the CS-TS pair. The orbit on the network-based side shown at the bottom of the image has been extended to

the entirety of the space of 1,536 network-based adinkras as shown in Fig. # 6 below taken from the work in [20].




Figure # 6: Orbits of Network-Based Adinkras

This current work has provided a tool that appears to be the most powerful in identifying network-based adinkras with 0-brane based supermultiplets. This tool is to impose matching conditions based on the representation space metrics $\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')]]$ and $\mathcal{G}[(\mathcal{R}), (\mathcal{R}')]]$ which leads to a projection operator.

(c.) Representation Space Metric Matching

The work of [18] fully enunciated an idea that when the two sets of equations in the forms of (32) and (33) are known, then it is possible to calculate (based on these) two respective fermionic holoraumy tensors and require their matching as the condition to when the network-based construction is the shadow of the 0-brane construction.

$$\mathcal{P}(\tilde{\mathcal{G}}[(\mathcal{R}), (\mathcal{R}')]]) = \mathcal{G}[(\mathcal{R}), (\mathcal{R}')]] \quad (35)$$

Thus, we believe these results constitute the essential elements of a proof that the kinematics of higher dimensional 4D, $\mathcal{N} = 1$ supersymmetric multiplets are embedded into the structure of network-based adinkras.

7 A Detailed Presentation On Holoraumy & Dimensional Reduction

In this chapter, we will take a detailed look at the interplay of dimensional reduction and the holoraumy tensor associated with multiplets in different dimensions

by looking at the examples of the 4D, $\mathcal{N} = 1$ vector supermultiplet and the 2D, $\mathcal{N} = 2$ twisted chiral multiplet to be contrasted with the simple reduction of the 4D, $\mathcal{N} = 1$ chiral supermultiplet.

7.1 Implementing Torus Compactification

We will naturally be led to a 2-manifold if we assume all the four dimensional fields do NOT depend on x^2 and x^3 . For the $\mathcal{N} = 1$ vector supermultiplet in 4D we thus have

$$\begin{aligned} D_a A_\mu &= (\gamma_\mu)_a{}^b \lambda_b \quad , \\ D_a \lambda_b &= -i \frac{1}{4} ([\gamma^\mu, \gamma^\nu]_{ab} (\partial_\mu A_\nu - \partial_\nu A_\mu) + (\gamma^5)_{ab} d) \quad , \\ D_a d &= i (\gamma^5 \gamma^\mu)_a{}^b \partial_\mu \lambda_b \quad , \end{aligned} \tag{36}$$

as our starting point. After a number of steps explained in detail in an appendix, we find

$$\begin{aligned} D_a \hat{A} &= (\gamma_2)_a{}^b \hat{\psi}_b \quad , \quad D_a \hat{B} = (\gamma_3)_a{}^b \hat{\psi}_b \quad , \\ D_a \hat{\psi}_b &= i (\gamma^2 \gamma^0)_{ab} (\partial_0 \hat{A}) + i (\gamma^2 \gamma^1)_{ab} (\partial_1 \hat{A}) \\ &\quad + i (\gamma^3 \gamma^0)_{ab} (\partial_0 \hat{B}) + i (\gamma^3 \gamma^1)_{ab} (\partial_1 \hat{B}) \\ &\quad - i (\gamma^0 \gamma^1)_{ab} \hat{F} + (\gamma^5)_{ab} \hat{G} \quad , \\ D_a \hat{F} &= (\gamma_1)_a{}^b (\partial_0 \hat{\psi}_b) - (\gamma_0)_a{}^b (\partial_1 \hat{\psi}_b) \quad , \\ D_a \hat{G} &= i (\gamma^5 \gamma^0)_a{}^b (\partial_0 \hat{\psi}_b) + i (\gamma^5 \gamma^1)_a{}^b (\partial_1 \hat{\psi}_b) \quad , \end{aligned} \tag{37}$$

and this is a formulation of the $\mathcal{N} = 2$ twisted chiral supermultiplet (TCS) in two dimensions. The process above is precisely the one that led in 1984 to the discovery of the twisted chiral supermultiplet [26].

On the other hand, it is simple to demand the fields of the 4D chiral supermultiplet

$$\begin{aligned} D_a A &= \psi_a \quad , \\ D_a B &= i (\gamma^5)_a{}^b \psi_b \quad , \\ D_a \psi_b &= i (\gamma^\mu)_{ab} (\partial_\mu A) - (\gamma^5 \gamma^\mu)_{ab} (\partial_\mu B) \\ &\quad - i C_{ab} F + (\gamma^5)_{ab} G \quad , \\ D_a F &= (\gamma^\mu)_a{}^b (\partial_\mu \psi_b) \quad , \\ D_a G &= i (\gamma^5 \gamma^\mu)_a{}^b (\partial_\mu \psi_b) \quad , \end{aligned} \tag{38}$$

should only depend on x^0 and x^1 yielding

$$\begin{aligned}
D_a A &= \psi_a , \\
D_a B &= i (\gamma^5)_a{}^b \psi_b , \\
D_a \psi_b &= i (\gamma^0)_{ab} (\partial_0 A) + i (\gamma^1)_{ab} (\partial_1 A) \\
&\quad - (\gamma^5 \gamma^0)_{ab} (\partial_0 B) - (\gamma^5 \gamma^1)_{ab} (\partial_1 B) \\
&\quad - i C_{ab} F + (\gamma^5)_{ab} G , \\
D_a F &= (\gamma^0)_a{}^b (\partial_0 \psi_b) + (\gamma^1)_a{}^b (\partial_1 \psi_b) , \\
D_a G &= i (\gamma^5 \gamma^0)_a{}^b (\partial_0 \psi_b) + i (\gamma^5 \gamma^1)_a{}^b (\partial_1 \psi_b) .
\end{aligned} \tag{39}$$

7.2 Reducing the 2D CS & TCS to 1D with Node Lowering

in order to find the covariant holonomy tensors associated with the two dimensional, $\mathcal{N} = 2$ chiral and twisted chiral supermultiplets, they each need to be reduced to one dimensional supermultiplets and node-lowering must be applied to their respective auxiliary fields. Once the respective holonomy tensors are found, then the metric on the space of these two representations can be calculated. The reduction of the CS and TCS to one dimension simply amounts to demanding all fields depend solely on x^0 .

Under this circumstance, we find for the CS

$$\begin{aligned}
D_a A &= \psi_a , \quad D_a B = i (\gamma^5)_a{}^b \psi_b , \\
D_a \psi_b &= i (\gamma^0)_{ab} (\partial_\tau A) - (\gamma^5 \gamma^0)_{ab} (\partial_\tau B) \\
&\quad - i C_{ab} F + (\gamma^5)_{ab} G , \\
D_a F &= (\gamma^0)_a{}^b (\partial_\tau \psi_b) , \\
D_a G &= i (\gamma^5 \gamma^0)_a{}^b (\partial_\tau \psi_b) ,
\end{aligned} \tag{40}$$

and we find for the TCS

$$\begin{aligned}
D_a \hat{A} &= (\gamma_2)_a{}^b \hat{\psi}_b , \quad D_a \hat{B} = (\gamma_3)_a{}^b \hat{\psi}_b , \\
D_a \hat{\psi}_b &= i (\gamma^2 \gamma^0)_{ab} (\partial_\tau \hat{A}) + i (\gamma^3 \gamma^0)_{ab} (\partial_\tau \hat{B}) \\
&\quad - i (\gamma^0 \gamma^1)_{ab} \hat{F} + (\gamma^5)_{ab} \hat{G} , \\
D_a \hat{F} &= (\gamma_1)_a{}^b (\partial_\tau \hat{\psi}_b) , \\
D_a \hat{G} &= i (\gamma^5 \gamma^0)_a{}^b (\partial_\tau \hat{\psi}_b) .
\end{aligned} \tag{41}$$

We introduce node lowering by implementing the re-definition of the “auxiliary fields” in each supermultiplet according to

$$F \rightarrow \partial_\tau F \quad , \quad G \rightarrow \partial_\tau G \quad , \quad \widehat{F} \rightarrow \partial_\tau \widehat{F} \quad , \quad \widehat{G} \rightarrow \partial_\tau \widehat{G} \quad , \quad (42)$$

so (40) becomes

$$\begin{aligned} D_a A &= \psi_a \quad , \quad D_a B = i(\gamma^5)_a{}^b \psi_b \quad , \\ D_a F &= (\gamma^0)_a{}^b \psi_b \quad , \quad D_a G = i(\gamma^5 \gamma^0)_a{}^b \psi_b \quad , \\ D_a \psi_b &= i(\gamma^0)_{ab} (\partial_\tau A) - (\gamma^5 \gamma^0)_{ab} (\partial_\tau B) \\ &\quad - i C_{ab} (\partial_\tau F) + (\gamma^5)_{ab} (\partial_\tau G) \quad , \end{aligned} \quad (43)$$

and for (41) we find

$$\begin{aligned} D_a \widehat{A} &= (\gamma_2)_a{}^b \widehat{\psi}_b \quad , \quad D_a \widehat{B} = (\gamma_3)_a{}^b \widehat{\psi}_b \quad , \\ D_a \widehat{F} &= (\gamma_1)_a{}^b \widehat{\psi}_b \quad , \quad D_a \widehat{G} = i(\gamma^5 \gamma^0)_a{}^b \widehat{\psi}_b \quad , \\ D_a \widehat{\psi}_b &= i(\gamma^2 \gamma^0)_{ab} (\partial_\tau \widehat{A}) + i(\gamma^3 \gamma^0)_{ab} (\partial_\tau \widehat{B}) \\ &\quad - i(\gamma^0 \gamma^1)_{ab} (\partial_\tau \widehat{F}) + (\gamma^5)_{ab} (\partial_\tau \widehat{G}) \quad . \end{aligned} \quad (44)$$

Given the equations in (44), the holonomy tensor associated with the 0-brane reduced 2D, $\mathcal{N} = 2$ twisted chiral supermultiplet can be calculated and is found to be the same as the 0-brane reduced 4D, $\mathcal{N} = 1$ vector supermultiplet as shown below

$$\begin{aligned} (\mathcal{F}_a{}^b{}^{(CS)})_c{}^d &= -(\gamma^5 \gamma^m)_a{}^b (\gamma^5 \gamma^0 \gamma_m)_c{}^d \quad , \\ (\mathcal{F}_a{}^b{}^{(TCS)})_c{}^d &= +\delta_a{}^b (\gamma^0)_c{}^d + (\gamma^5)_a{}^b (\gamma^5 \gamma^0)_c{}^d + (\gamma^5 \gamma^0)_a{}^b (\gamma^5)_c{}^d \quad . \end{aligned} \quad (45)$$

Now a few calculations show

$$\begin{aligned} (\mathcal{F}_{ab}{}^{(CS)})_e{}^f (\mathcal{F}_{cd}{}^{(TCS)})_f{}^h - (\mathcal{F}_{cd}{}^{(TCS)})_e{}^f (\mathcal{F}_{ab}{}^{(CS)})_f{}^h &= 0 \quad , \\ (\mathcal{F}_{ab}{}^{(CS)})_e{}^f (\mathcal{F}_{cd}{}^{(TCS)})_f{}^e &= 0 \quad . \end{aligned} \quad (46)$$

The holonomy tensors of the 2D, $\mathcal{N} = 2$ chiral supermultiplet are found to commute with those of the 2D, $\mathcal{N} = 2$ twisted chiral supermultiplet. The final equation in (46) implies starting from these two distinct 2D, $\mathcal{N} = 2$ chiral supermultiplets are “orthogonal,” i.e.

$$\widetilde{\mathcal{G}}[(CS), (TCS)] = 0 \quad , \quad (47)$$

just as the 4D, $\mathcal{N} = 1$ chiral supermultiplet is “orthogonal” to the 4D, $\mathcal{N} = 1$ vector supermultiplet.

8 How Ignoring SUSY Holography Can Lead To Violation of Lorentz Symmetry In Uplifting

Let us begin by considering the two adinkra graphs shown below.




Figure # 7: Two valise adinkra networks with opposite red-parity

It is seen they are identical but with one exception. All the solid red links in Adnk-A are replaced by dashed red links in Adnk-B and vice versa.

Now we can use the adinkra in (Adnk-A) to find the associated L-matrices (and R-matrices which are simply the transposed versions of the L-matrices). After such explicit expressions are found (see Appendix E), the corresponding $\tilde{V}$ matrices are calculated. The results of these calculations further imply

$$\begin{aligned} \tilde{V}_{12}^{(\text{Adnk-A})} &= \tilde{V}_{34}^{(\text{Adnk-A})} , & \tilde{V}_{23}^{(\text{Adnk-A})} &= - \tilde{V}_{14}^{(\text{Adnk-A})} , \\ \tilde{V}_{13}^{(\text{Adnk-A})} &= \tilde{V}_{24}^{(\text{Adnk-A})} . \end{aligned} \quad (48)$$

Similarly,

$$\begin{aligned} \tilde{V}_{12}^{(\text{Adnk-B})} &= - \tilde{V}_{34}^{(\text{Adnk-B})} , & \tilde{V}_{23}^{(\text{Adnk-B})} &= \tilde{V}_{14}^{(\text{Adnk-B})} , \\ \tilde{V}_{13}^{(\text{Adnk-B})} &= - \tilde{V}_{24}^{(\text{Adnk-B})} . \end{aligned} \quad (49)$$

Furthermore, these same calculations make it clear

$$\begin{aligned} \tilde{V}_{12}^{(\text{Adnk-A})} &= - \tilde{V}_{12}^{(\text{Adnk-B})} , & \tilde{V}_{13}^{(\text{Adnk-A})} &= - \tilde{V}_{13}^{(\text{Adnk-B})} , \\ \tilde{V}_{14}^{(\text{Adnk-A})} &= \tilde{V}_{14}^{(\text{Adnk-B})} , \\ \tilde{V}_{IJ}^{(\text{Adnk-A})} \tilde{V}_{KL}^{(\text{Adnk-B})} - \tilde{V}_{KL}^{(\text{Adnk-B})} \tilde{V}_{IJ}^{(\text{Adnk-A})} &\neq 0 . \end{aligned} \quad (50)$$

If one makes the claim either (Adnk-A) or (Adnk-B) is the 0-brane reduction of the chiral supermultiplet and the other is the 0-brane reduction of the twisted chiral multiplet, then one is forced to one of two conclusions:

- (a.) the assignment of these adinkra as ‘shadows’ of the 2D, $\mathcal{N} = 2$ supermultiplets is inconsistent, or

(b.) SUSY holography is violated.

The reason one is driven to these is we have previously argued the commutator of the $\tilde{V}$ -matrices is tantamount to the evaluation of the fermionic holoraumy tensor on the supermultiplet via SUSY holography. Via the first equation of (46) and SUSY holography we conclude the commutator in (50) should be zero.

In Appendix C, it is shown how the original derivation of the 2D, $\mathcal{N} = 2$ twisted chiral supermultiplet is obtained from the 4D, $\mathcal{N} = 1$ vector supermultiplet in the real basis used in this work. We now wish to build on this derivation to show that applying the traditional understanding of how the 2D, $\mathcal{N} = 2$ twisted chiral supermultiplet (CS) is related to the 4D, $\mathcal{N} = 1$ vector supermultiplet (VS) together with the assertion that one of the adinkras shown in (Adnk-A) and (Adnk-B) corresponds to the CS and the other to the VS means implies a violation of Lorentz invariance.

The closure of the off-shell (and on-shell) algebra is not dependent on the choice of γ -matrices used. It is only their algebraic properties that are important. Thus the calculation of the holoraumy on the 0-brane side has the same form when expressed in terms of γ -matrices. The calculation of the holoraumy on the adinkra side will depend on the V and $\tilde{V}$ matrices of the adinkra used.

So, if one claims the chiral supermultiplet has (Adnk-A) as its adinkra shadow, the projection of the 0-brane-based holoraumy tensor onto the adinkra-based holoraumy tensor implies there must exist constants $\mathcal{S}_1^{ij}$, $\mathcal{S}_2^{ij}$, and $\mathcal{S}_3^{ij}$ so

$$\begin{aligned}\gamma^{12} &= \mathcal{S}_1^{12} \tilde{V}_{12}^{(\text{Adnk-A})} + \mathcal{S}_2^{12} \tilde{V}_{13}^{(\text{Adnk-A})} + \mathcal{S}_3^{12} \tilde{V}_{14}^{(\text{Adnk-A})} , \\ \gamma^{23} &= \mathcal{S}_1^{23} \tilde{V}_{12}^{(\text{Adnk-A})} + \mathcal{S}_2^{23} \tilde{V}_{13}^{(\text{Adnk-A})} + \mathcal{S}_3^{23} \tilde{V}_{14}^{(\text{Adnk-A})} , \\ \gamma^{31} &= \mathcal{S}_1^{31} \tilde{V}_{12}^{(\text{Adnk-A})} + i \mathcal{S}_2^{31} \tilde{V}_{13}^{(\text{Adnk-A})} + \mathcal{S}_3^{31} \tilde{V}_{14}^{(\text{Adnk-A})} .\end{aligned}\tag{51}$$

and this equation defines an explicit representation.

If it is asserted the adinkra in (Adnk-B) is the valise shadow of the 0-brane reduced valise version of the vector supermultiplet shown in (3.), the holoraumy tensor of (13) on the adinkra side must be related to the holoraumy tensor in (4) that originates from the VS supermultiplet. In that case there must exist constants $\mathcal{S}_1, \dots, \mathcal{S}_9$ such that

$$\begin{aligned}\gamma^0 &= \mathcal{S}_1 \tilde{V}_{12}^{(\text{Adnk-B})} + i \mathcal{S}_2 \tilde{V}_{13}^{(\text{Adnk-B})} + \mathcal{S}_3 \tilde{V}_{14}^{(\text{Adnk-B})} , \\ \gamma^0 \gamma^5 &= \mathcal{S}_4 \tilde{V}_{12}^{(\text{Adnk-B})} + \mathcal{S}_5 \tilde{V}_{13}^{(\text{Adnk-B})} + \mathcal{S}_6 \tilde{V}_{14}^{(\text{Adnk-B})} , \\ \gamma^5 &= \mathcal{S}_7 \tilde{V}_{12}^{(\text{Adnk-B})} + \mathcal{S}_8 \tilde{V}_{13}^{(\text{Adnk-B})} + \mathcal{S}_9 \tilde{V}_{14}^{(\text{Adnk-B})} .\end{aligned}\tag{52}$$

Furthermore, since no explicit representation of the γ -matrices need be chosen, this equation defines an explicit representation.

The freedom in choosing the $\mathcal{S}$ -coefficients is not as large as it would first appear. The commutator algebra of the γ matrices can easily be derived from (5). On the other hand, the commutation relations of the $\tilde{V}$ matrices are determined from the adinkras and their adjacency matrices.

Combining the results in (8), (51), and (52) leads to a major problem. Once one determines a choice of the $\mathcal{S}$ coefficients, then one can calculate the traces

$$\begin{aligned} t^{0\ ij} &= \text{Tr} [\gamma^0 \gamma^{ij}] \quad , \\ t^{5\ ij} &= \text{Tr} [\gamma^5 \gamma^{ij}] \quad , \\ t^{05\ ij} &= \text{Tr} [\gamma^0 \gamma^5 \gamma^{ij}] \quad . \end{aligned} \tag{53}$$

For any proper choice of γ -matrices all of these traces vanish. Also for any proper choice of γ -matrices we have

$$[\gamma^0, \gamma^{ij}] = 0 \quad , \quad [\gamma^5, \gamma^{ij}] = 0 \quad , \quad [\gamma^0 \gamma^5, \gamma^{ij}] = 0 \quad . \tag{54}$$

However, when one

- (a.) uses the choice of adinkras shown in (Adnk-A) and in (Adnk-B),
- (b.) asserts these are the valise adinkra shadows of the chiral supermultiplet and the vector supermultiplet,
- (c.) implies the set $\{\tilde{V}_{12}^{(\text{Adnk-A})}, \tilde{V}_{13}^{(\text{Adnk-A})}, \tilde{V}_{14}^{(\text{Adnk-A})}\}$ becomes linearly dependent on the set $\{\tilde{V}_{12}^{(\text{Adnk-B})}, \tilde{V}_{13}^{(\text{Adnk-B})}, \tilde{V}_{14}^{(\text{Adnk-B})}\}$ as shown in (8),
- (d.) forces the set $\{\gamma^{ij}\}$ to be linearly dependent on the set $\{\gamma^0, \gamma^0 \gamma^5, \gamma^5\}$
- (e.) leads to the non-vanishing of the traces in (53), and
- (f.) ends by violating the conditions in (54).

Lorentz invariance is violated! How can all this be avoided?

If one repeats all the calculations based on the network-based adinkras in Fig. # 7 (see also Appendix E), all the problems disappear. Explicitly and by using the same rainbow assignment. We find the results

$$\begin{aligned} \tilde{V}_{12}^{(\text{Adnk-C})} &= -\tilde{V}_{34}^{(\text{Adnk-C})} \quad , \quad \tilde{V}_{23}^{(\text{Adnk-C})} = -\tilde{V}_{14}^{(\text{Adnk-C})} \quad , \\ \tilde{V}_{13}^{(\text{Adnk-C})} &= \tilde{V}_{24}^{(\text{Adnk-C})} \quad , \\ \tilde{V}_{12}^{(\text{Adnk-D})} &= \tilde{V}_{34}^{(\text{Adnk-D})} \quad , \quad \tilde{V}_{23}^{(\text{Adnk-D})} = \tilde{V}_{14}^{(\text{Adnk-D})} \quad , \\ \tilde{V}_{13}^{(\text{Adnk-D})} &= -\tilde{V}_{24}^{(\text{Adnk-D})} \quad . \end{aligned} \tag{55}$$




Figure # 8: Two valise adinkra networks

The $\tilde{V}$ matrices associated with the (Adnk-C) are linearly independent of the $\tilde{V}$ matrices associated with the (Adnk-D). Further it is possible to derive

$$\tilde{V}_{IJ}^{(\text{Adnk-C})} \tilde{V}_{KL}^{(\text{Adnk-D})} - \tilde{V}_{KL}^{(\text{Adnk-D})} \tilde{V}_{IJ}^{(\text{Adnk-C})} = 0 \quad . \quad (56)$$

which shows SUSY holography is restored. It is also the case

$$\text{Tr} \left[\tilde{V}_{IJ}^{(\text{Adnk-C})} \tilde{V}_{KL}^{(\text{Adnk-D})} \right] = 0 \quad , \quad (57)$$

which are the necessary and sufficient conditions to avoid the violations of Lorentz symmetry.

We close this discussion by noting a necessary condition for two adinkras to represent the shadows of two distinct higher dimensional supermultiplets is that their respective L-matrices must be linearly independent. One way to do this is to choose the permutation cycles related to the two quartets of L-matrices to lie in different sectors of the Venn diagram shown in (Fig. 2) of [20]. If the shadows of two distinct higher dimensional supermultiplets possess the same permutation cycles, then a proper choice of Boolean factors can also be used to maintain linear independence.

9 Holoraumy Tensors for the Valise Real Scalar Supermultiplet

The transformation laws for the fields of the real scalar supermultiplet in valise form are given by the equations

$$\begin{aligned} D_a K &= \zeta_a \\ D_a M &= \frac{1}{2} \Lambda_a - \frac{1}{2} (\gamma^0)_a^d \zeta_d \\ D_a N &= -i \frac{1}{2} (\gamma^5)_a^d \Lambda_d + i \frac{1}{2} (\gamma^5 \gamma^0)_a^d \zeta_d \\ D_a U_0 &= i \frac{1}{2} (\gamma^5 \gamma_0)_a^d \Lambda_d - i \frac{1}{2} (\gamma^5)_a^d \zeta_d \end{aligned}$$

$$\begin{aligned}
D_a U_m &= i \frac{1}{2} (\gamma^5 \gamma_m)_a^d \Lambda_d - i \frac{1}{2} (\gamma^5 \gamma^0 \gamma_m)_a^d \zeta_d \\
D_a d &= - (\gamma^0)_a^d \Lambda_d \\
D_a \zeta_b &= i (\gamma^0)_{ab} \partial_\tau K + (\gamma^5 \gamma^\mu)_{ab} \partial_\tau U_\mu + i C_{ab} \partial_\tau M + (\gamma^5)_{ab} \partial_\tau N \\
D_a \Lambda_b &= i (\gamma^0)_{ab} \partial_\tau M + (\gamma^5 \gamma^0)_{ab} \partial_\tau N + (\gamma^5 \gamma^0 \gamma^\nu)_{ab} \partial_\tau U_\nu + i C_{ab} \partial_\tau d
\end{aligned} \tag{58}$$

and to find bosonic holoraumy tensor for the real scalar supermultiplet we note

$$\begin{aligned}
[D_a, D_b] K &= i 2 C_{ab} \partial_\tau M + 2 (\gamma^5)_{ab} \partial_\tau N + 2 (\gamma^5 \gamma^0)_{ab} \partial_\tau U_0 + 2 (\gamma^5 \gamma^m)_{ab} \partial_\tau U_m \\
[D_a, D_b] M &= 2 (\gamma^5 \gamma^0)_{ab} \partial_\tau N - 2 (\gamma^5)_{ab} \partial_\tau U_0 + i C_{ab} \partial_\tau (d - K) \\
[D_a, D_b] N &= -2 (\gamma^5 \gamma^0)_{ab} \partial_\tau M + i 2 C_{ab} \partial_\tau U_0 + (\gamma^5)_{ab} \partial_\tau (d - K) \\
[D_a, D_b] U_0 &= 2 (\gamma^5)_{ab} \partial_\tau M - i 2 C_{ab} \partial_\tau N + (\gamma^5 \gamma^0)_{ab} \partial_\tau (d - K) \\
[D_a, D_b] U_m &= - (\gamma^5 \gamma_m)_{ab} \partial_\tau (d + K) - i (\gamma^0 [\gamma_m, \gamma^n])_{ab} \partial_\tau U_n \\
[D_a, D_b] d &= -2 i C_{ab} \partial_\tau M - 2 (\gamma^5)_{ab} \partial_\tau N - 2 (\gamma^5 \gamma^0)_{ab} \partial_\tau U_0 + 2 (\gamma^5 \gamma^m)_{ab} \partial_\tau U_m
\end{aligned} \tag{59}$$

It is clear the linear combinations $(d \pm K)$ possess some significance suggesting these be rewritten in the forms

$$\begin{aligned}
[D_a, D_b] (d + K) &= 4 (\gamma^5 \gamma^m)_{ab} \partial_\tau U_m \\
[D_a, D_b] M &= 2 (\gamma^5 \gamma^0)_{ab} \partial_\tau N - 2 (\gamma^5)_{ab} \partial_\tau U_0 + i C_{ab} \partial_\tau (d - K) \\
[D_a, D_b] N &= -2 (\gamma^5 \gamma^0)_{ab} \partial_\tau M + i 2 C_{ab} \partial_\tau U_0 + (\gamma^5)_{ab} \partial_\tau (d - K) \\
[D_a, D_b] U_0 &= 2 (\gamma^5)_{ab} \partial_\tau M - i 2 C_{ab} \partial_\tau N + (\gamma^5 \gamma^0)_{ab} \partial_\tau (d - K) \\
[D_a, D_b] U_m &= - (\gamma^5 \gamma_m)_{ab} \partial_\tau (d + K) - i (\gamma^0 [\gamma_m, \gamma^n])_{ab} \partial_\tau U_n \\
[D_a, D_b] (d - K) &= -4 i C_{ab} \partial_\tau M - 4 (\gamma^5)_{ab} \partial_\tau N - 4 (\gamma^5 \gamma^0)_{ab} \partial_\tau U_0
\end{aligned} \tag{60}$$

These equations are interesting as they clearly identify two distinct subsets

$$subset_1 = \{ (d + K), U_m \} \quad , \quad subset_2 = \{ (d - K), M, N, U_0 \} \tag{61}$$

Calculations for the fermionic holoraumy tensor for the real scalar supermultiplet yield

$$\begin{aligned}
[D_a, D_b] \zeta_c &= i C_{ab} (\gamma^0)_c^d \partial_\tau [\zeta_d - (\gamma^0)_d^e \Lambda_e] \\
&\quad - i (\gamma^5)_{ab} (\gamma^5 \gamma^0)_c^d \partial_\tau [\zeta_d - (\gamma^0)_d^e \Lambda_e] \\
&\quad + i (\gamma^5 \gamma^0)_{ab} (\gamma^5)_c^d \partial_\tau [\zeta_d - (\gamma^0)_d^e \Lambda_e] \\
&\quad + i (\gamma^5 \gamma^m)_{ab} (\gamma^5 \gamma^0 \gamma_m)_c^d \partial_\tau [\zeta_d + (\gamma^0)_d^e \Lambda_d] \\
[D_a, D_b] \Lambda_c &= i C_{ab} (\gamma^0)_c^d \partial_\tau [(\gamma^0)_d^e (\zeta_e - (\gamma^0)_e^f \Lambda_f)] \\
&\quad + i (\gamma^5)_{ab} (\gamma^5 \gamma^0)_c^d \partial_\tau [(\gamma^0)_d^e (\zeta_e - (\gamma^0)_e^f \Lambda_f)] \\
&\quad - i (\gamma^5 \gamma^0)_{ab} (\gamma^5)_c^d \partial_\tau [(\gamma^0)_d^e (\zeta_e - (\gamma^0)_e^f \Lambda_f)] \\
&\quad - i (\gamma^5 \gamma^m)_{ab} (\gamma^5 \gamma^0 \gamma_m)_c^d \partial_\tau [(\gamma^0)_d^e (\zeta_e + (\gamma^0)_e^f \Lambda_f)]
\end{aligned} \tag{62}$$

or alternatively

$$\begin{aligned}
[D_a, D_b] [\zeta_c + (\gamma^0)_c{}^d \Lambda_d] &= +i2(\gamma^5 \gamma^m)_{ab} (\gamma^5 \gamma^0 \gamma_m)_c{}^d \partial_\tau [\zeta_d + (\gamma^0)_d{}^e \Lambda_e] \\
[D_a, D_b] [\zeta_c - (\gamma^0)_c{}^d \Lambda_d] &= -i2C_{ab} (\gamma^0)_c{}^d \partial_\tau \left[(\zeta_d - (\gamma^0)_d{}^f \Lambda_f) \right] \\
&\quad + i2(\gamma^5)_{ab} (\gamma^5 \gamma^0)_c{}^d \partial_\tau \left[(\zeta_d - (\gamma^0)_d{}^f \Lambda_f) \right] \\
&\quad + i2(\gamma^5 \gamma^0)_{ab} (\gamma^5)_c{}^d \partial_\tau \left[(\zeta_d - (\gamma^0)_d{}^f \Lambda_f) \right]
\end{aligned} \tag{63}$$

We observe

$$D_a(d \pm K) = -[(\gamma^0)_a{}^b \Lambda_b \mp \zeta_a] \tag{64}$$

along with the expressions for the fermionic holoraumy tensor above, this suggests the definitions

$$\begin{aligned}
\Phi_\Lambda &\equiv ((d - K), M, N, U_0 | U_m, (d + K)) \quad , \\
\Psi_{\hat{\Lambda}} &\equiv (\zeta_a + (\gamma^0)_a{}^b \Lambda_b | \zeta_a - (\gamma^0)_a{}^b \Lambda_b) \quad ,
\end{aligned} \tag{65}$$

and we combine these equations to find,

$$[D_a, D_b] \Phi_\Lambda = -i2(\mathcal{B}_{ab})_\Lambda{}^\Delta \partial_\tau \Phi_\Delta \tag{66}$$

$$[D_a, D_b] \Psi_{\hat{\Lambda}} = -i2(\mathcal{F}_{ab})_{\hat{\Lambda}}{}^{\hat{\Delta}} \partial_\tau \Psi_{\hat{\Delta}} \tag{67}$$

for the explicit representation-dependent forms of $(\mathcal{B}_{ab})_\Lambda{}^\Delta$ and $(\mathcal{F}_{ab})_{\hat{\Lambda}}{}^{\hat{\Delta}}$. The basis for Φ_Λ and $\Psi_{\hat{\Lambda}}$ has the property that both $(\mathcal{B}_{ab})_\Lambda{}^\Delta$ and $(\mathcal{F}_{ab})_{\hat{\Lambda}}{}^{\hat{\Delta}}$ are block diagonal with 4×4 matrices on the main diagonal of the 8×8 matrices. This completes the derivation of the holoraumy tensors on the 0-brane side of the correspondence.

The splitting of the real scalar supermultiplet into the two distinct subsets seen in (65) can be illustrated⁵ as below using adinkras.




Figure # 9: Illustration of Diagonalized Holoraumy

The subset of fields shown to the left of the “vertical bars” in (65) constitute the fields seen in the middle adinkra of Fig. # 9 while subset of fields shown to the right

⁵This image appeared in “Symbols of Power” in [32].

of the “vertical bars” constitute the fields seen in the rightmost adinkra. The leftmost adinkra corresponds to the real scalar supermultiplet. It can be seen by the equations in (63), the splitting is accomplished by choosing between the basis defined by the linear combination of fermions defined by $[\zeta_c \pm (\gamma^0)_c{}^d \Lambda_d]$ as these linear combinations control how the fermions in the original real scalar superfield must be rearranged to obtain the fermions in the sub-multiplets. On the network-based adinkra side, this same splitting is accomplished by the use of error-correcting codes [3, 4, 5]. Thus we find the result the use of error-correcting codes on the network-based adinkra side corresponds to the splitting of $SO(1,3)$ fermions into covariant $SO(3)$ ones on the 0-brane reduced side.

We believe this is a very telling sign of the linkage between 1D, N -extended SUSY together with error-correcting codes and the emergence of Lorentz symmetry in the higher dimensional space-time manifolds.

On the adinkra side, we find the following results. We define the adinkra node “vectors” via

$$\begin{aligned}\Phi_i &= (d, M + U_2, U_0 + U_1, N + U_3, -N + U_3, U_0 + U_1, -M + U_2, K) \quad , \\ \Psi_{\hat{k}} &= (\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, \zeta_1, \zeta_2, \zeta_3, \zeta_4) \quad ,\end{aligned}\tag{68}$$

and using this definition we find the L-matrices and R-matrices are given by

$$\begin{aligned}L_1 &= (128)_b\{(12)(38)(57)\} \quad , \quad L_2 = (22)_b\{(34)(53)(67)\} \quad , \\ L_3 &= (42)_b\{(14)(28)(31)\} \quad , \quad L_4 = (230)_b\{(13)(27)(46)(51)\} \quad ,\end{aligned}\tag{69}$$

$$\begin{aligned}R_1 &= (64)_b\{(12)(36)(58)\} \quad , \quad R_2 = (11)_b\{(35)(43)(68)\} \quad , \\ R_3 &= (168)_b\{(13)(24)(78)\} \quad , \quad R_4 = (122)_b\{(15)(26)(31)(47)\} \quad ,\end{aligned}\tag{70}$$

For the V -matrices and $\tilde{V}$ -matrices we find,

$$V_{IJ} = \frac{1}{2} (L_I R_J - L_J R_I) \quad , \quad \tilde{V}_{IJ} = \frac{1}{2} (R_I L_J - R_J L_I) \quad ,\tag{71}$$

and enumerate the non-vanishing and non-redundant values:

$$\begin{aligned}V_{12} &= (149)_b\{(12)(34)(56)(78)\} \\ V_{13} &= (163)_b\{(14)(23)(58)(67)\} \\ V_{14} &= (184)_b\{(16)(25)(38)(47)\} \\ V_{23} &= (57)_b\{(13)(24)(57)(68)\} \\ V_{24} &= (30)_b\{(15)(26)(37)(48)\} \\ V_{34} &= (141)_b\{(17)(28)(35)(46)\}\end{aligned}\tag{72}$$

$$\begin{aligned}
\tilde{V}_{12} &= (46)_b \{ (12)(37)(48)(56) \} \\
\tilde{V}_{13} &= (226)_b \{ (18)(24)(36)(57) \} \\
\tilde{V}_{14} &= (165)_b \{ (17)(23)(46)(58) \} \\
\tilde{V}_{23} &= (204)_b \{ (14)(28)(35)(67) \} \\
\tilde{V}_{24} &= (139)_b \{ (13)(27)(45)(68) \} \\
\tilde{V}_{34} &= (184)_b \{ (15)(26)(34)(78) \} \quad .
\end{aligned} \tag{73}$$

For the algebra of the V -matrices and $\tilde{V}$ -matrices we find,

$$\frac{1}{2}[V_{IJ}, V_{KL}] = -\delta_{IK}V_{JL} + \delta_{IL}V_{JK} + \delta_{JK}V_{IL} - \delta_{JL}V_{IK} \quad , \tag{74}$$

$$\frac{1}{2}[\tilde{V}_{IJ}, \tilde{V}_{KL}] = -\delta_{IK}\tilde{V}_{JL} + \delta_{IL}\tilde{V}_{JK} + \delta_{JK}\tilde{V}_{IL} - \delta_{JL}\tilde{V}_{IK} \quad , \tag{75}$$

and the factors of one-half appearing in these informs us these constitute spinor representations of $SO(4)$. We can form the following linear combinations of the $\tilde{V}$ -matrices

$$\tilde{V}_{IJ}^{(\pm)} = \frac{1}{2} \left[\tilde{V}_{IJ} \pm \frac{1}{2} \epsilon_{IJKL} \tilde{V}_{KL} \right] \tag{76}$$

and due to (74) we find

$$\left[\tilde{V}_{IJ}^{(\pm)} , \tilde{V}_{KL}^{(\mp)} \right] = 0 \tag{77}$$

It is now of interest to note the possibility to define two sets of matrices denoted by $[\mathcal{S}_{\text{IJK}}^{(\pm)}]$ and defined by the expressions

$$V_{JK}^{(\pm)} L_{\text{I}} - L_{\text{I}} \tilde{V}_{JK}^{(\pm)} = [\mathcal{S}_{\text{IJK}}^{(\pm)}] \quad , \tag{78}$$

since the equivalent expression of the 0-brane side contain information about spin and R-symmetry spin of the fields in the real scalar superfield. We plan to return to this topic in a future effort.

10 Extending Holoraumy to 4D, $\mathcal{N} > 1$ SUSY

All our previous discussion of holoraumy has been within the arena of four dimensional (or less) simple supersymmetry. In this section, we outline the general extension of these ideas to theories with more supercharges.

We begin by noting a simple counting argument. In order to describe an extended supersymmetry in spacetime, we replace the supercovariant derivative via the rule

$$D_a \rightarrow D_{ai} \quad , \tag{79}$$

where the i subscript counts the number of four dimensional spacetime supercharges. It thus ranges as $1, \dots, \mathcal{N}$, where now $\mathcal{N}$ can be any integer. The algebra of this extended SUSY covariant derivative is simply

$$\{D_{ai}, D_{bj}\} = i 2 \delta_{ij} (\gamma^\mu)_{ab} \partial_\mu \quad , \quad (80)$$

for all *off-shell* SUSY representations. Via “0-brane reduction” [21] and “node lowering” all *off-shell* representations of this algebra can be expressed in such a way that this anti-commutation relationship becomes

$$\{D_{ai}, D_{bj}\} = i 2 \delta_{ij} (\gamma^0)_{ab} \partial_\tau \quad . \quad (81)$$

In many places in the SUSY literature, it is customary to deform the anti-commutator relation in (80) so it reads

$$\begin{aligned} \{D_{ai}, D_{bj}\} = & i 2 \delta_{ij} (\gamma^\mu)_{ab} \partial_\mu + (\Delta_{ai bj}^{[1]}) \mathcal{Z}_{[1]} + (\Delta_{ai bj}^{[2]}) \mathcal{Z}_{[2]} \\ & + (\Delta_{ai bj}^{[3]}) \mathcal{Z}_{[3]} + (\Delta_{ai bj}^{[4]}) \mathcal{Z}_{[4]} + \dots \quad , \end{aligned} \quad (82)$$

where the operators $\mathcal{Z}_{[1]}, \mathcal{Z}_{[2]}, \mathcal{Z}_{[3]}, \mathcal{Z}_{[4]}$, etc. are called “off-shell” central charges if they vanish when the equations of motion, derivable from some Lagrangian, are imposed. The quantities $(\Delta_{ai bj}^{[1]})$, etc. are simply some set of constants. Although we do not have a general proof about the behavior of “off-shell” central charges under 0-brane reduction, we do have at least one example which suggests something interesting about the result one can expect generally.

In the work of [28], the Wess-Fayet hypermultiplet [29] (which has widely been used to represent $\mathcal{N} = 2$ supermatter), was studied under 0-brane reduction and “node lowering.” The result of the study indicated in this limit, one finds

$$\mathcal{Z}_{[1]} = z_{[1]} \partial_\tau \quad , \quad \mathcal{Z}_{[2]} = z_{[2]} \partial_\tau \quad , \quad \dots \quad , \quad (83)$$

for some constants $z_{[1]}, z_{[2]}, z_{[3]}, \dots$ on this supermultiplet. In the following, we make the *assumption* this behavior is valid for *all* theories which possess off-shell central charges. Under this assumption, when (82) is subjected to 0-brane reduction and node-lowering we expect the equation

$$\begin{aligned} \{D_{ai}, D_{bj}\} = & i 2 \delta_{ij} (\gamma^0)_{ab} \partial_\tau + (\Delta_{ai bj}^{[1]}) z_{[1]} \partial_\tau + (\Delta_{ai bj}^{[2]}) z_{[2]} \partial_\tau \\ & + (\Delta_{ai bj}^{[3]}) z_{[3]} \partial_\tau + (\Delta_{ai bj}^{[4]}) z_{[4]} \partial_\tau + \dots \quad , \end{aligned} \quad (84)$$

to emerge. In the case of the hypermultiplet, we also found the conditions

$$\begin{aligned} \delta^{ij} (\gamma^\mu)^{ab} (\Delta_{ai bj}^{[1]}) &= \delta^{ij} (\gamma^\mu)^{ab} (\Delta_{ai bj}^{[2]}) = \delta^{ij} (\gamma^\mu)^{ab} (\Delta_{ai bj}^{[3]}) = \\ \delta^{ij} (\gamma^\mu)^{ab} (\Delta_{ai bj}^{[4]}) &= \dots = 0 \quad . \end{aligned} \quad (85)$$

Once more we assert this is also the case for any general 0-brane reduction and node-lowering on any SUSY representation. The result in (85) and the fact for all the Δ -symbols we have the identity

$$(\Delta_{ai bj}^{\widehat{[1]}}) = (\Delta_{bj ai}^{\widehat{[1]}}) , \quad (86)$$

implies we can count their linearly independent parts easily. Since these quantities are bilinear in spinor indices a and b they can be expanded over the six elements

$$C_{ab} , \quad (\gamma^5)_{ab} , \quad (\gamma^5 \gamma^\mu)_{ab} , \quad (87)$$

(which are antisymmetric on the exchange of the a and b indices) and the ten elements

$$(\gamma^\mu)_{ab} , \quad (\gamma^\mu \wedge \gamma^\nu)_{ab} , \quad (88)$$

which are symmetric on the exchange of the a and b . In order to realize the symmetry as in (86) we must pair the six elements in (87) with an antisymmetric tensor $\mathcal{A}_{ij}$ to form the objects

$$C_{ab} \mathcal{A}_{ij} , \quad (\gamma^5)_{ab} \mathcal{A}_{ij} , \quad (\gamma^5 \gamma^\mu)_{ab} \mathcal{A}_{ij} , \quad (89)$$

which gives a total of $6 \times \mathcal{N}(\mathcal{N}-1)/2$ tensors. In a similar manner in order to realize the symmetry as in (86) we must pair the 10 elements in (88) with an symmetric tensor $\mathcal{S}_{ij}$ to form the objects

$$(\gamma^\mu)_{ab} \mathcal{S}_{ij} , \quad (\gamma^\mu \wedge \gamma^\nu)_{ab} \mathcal{S}_{ij} , \quad (90)$$

which gives a total of $10 \times \mathcal{N}(\mathcal{N}+1)/2$ tensors. We also note the objects in (90) can be replaced by

$$(\gamma^\mu)_{ab} \delta_{ij} , \quad (\gamma^\mu)_{ab} \mathcal{S}_{ij}^{(T)} , \quad (\gamma^\mu \wedge \gamma^\nu)_{ab} \mathcal{S}_{ij} , \quad (91)$$

where $\delta^{ij} \mathcal{S}_{ij}^{(T)} = 0$. Thus, without loss of generality, the Δ -tensors appearing in (84) - (86) can be expanded in the product tensors in (89) and (91).

Note two identities

$$\binom{d_F}{2} \binom{\mathcal{N}}{2} + \binom{d_F+1}{2} \binom{\mathcal{N}+1}{2} = \binom{d_F \mathcal{N} + 1}{2} , \quad (92)$$

$$\binom{d_F+1}{2} \binom{\mathcal{N}}{2} + \binom{d_F}{2} \binom{\mathcal{N}+1}{2} = \binom{d_F \mathcal{N}}{2} , \quad (93)$$

Adding together the number of independent constant tensors in (89) and (90) gives

$$\binom{4}{2} \binom{\mathcal{N}}{2} + \binom{5}{2} \binom{\mathcal{N}+1}{2} = \binom{4\mathcal{N}+1}{2} . \quad (94)$$

This is a special case of the identity in (92) for $d_F = 4$. On the other hand, if we interchange the factors of 6 and 10 in (94), we obtain

$$\binom{5}{2} \binom{\mathcal{N}}{2} + \binom{4}{2} \binom{\mathcal{N}+1}{2} = \binom{4\mathcal{N}}{2} , \quad (95)$$

and this is a special case of the identity in (93) for $d_F = 4$. The result in (95) is indicative the objects

$$\begin{aligned} & (\gamma^\mu)_{ab} \mathcal{A}_{ij} , \quad (\gamma^\mu \wedge \gamma^\nu)_{ab} \mathcal{A}_{ij} , \\ & C_{ab} \delta_{ij} , \quad (\gamma^5)_{ab} \delta_{ij} , \quad (\gamma^5 \gamma^\mu)_{ab} \delta_{ij} , \\ & C_{ab} \mathcal{S}_{ij}^{(T)} , \quad (\gamma^5)_{ab} \mathcal{S}_{ij}^{(T)} , \quad (\gamma^5 \gamma^\mu)_{ab} \mathcal{S}_{ij}^{(T)} , \end{aligned} \quad (96)$$

are all anti-symmetrical under the exchange of the pair of indices ai with bj . They are antisymmetric and their number is given by the RHS in (95) indicates these are in the defining representation of the group $\text{SO}(4\mathcal{N})$.

The operator equation

$$[D_{ai} , D_{bj}] = (\mathcal{H}_{aibj}) , \quad (97)$$

implies the matrices in (96) must form a representation of the generators of $\text{SO}(4\mathcal{N})$. The RHS of (97) can be decomposed in the basis in (??) and re-expressed as

$$\begin{aligned} [D_{ai} , D_{bj}] &= i(\gamma^\mu)_{ab} (\mathcal{A}_{ij} \cdot \mathcal{H}_\mu) + i(\gamma^\mu \wedge \gamma^\nu)_{ab} (\mathcal{A}_{ij} \cdot \mathcal{H}_{\mu\nu}) \\ &+ iC_{ab} (\mathcal{S}_{ij} \cdot \mathcal{H}) + (\gamma^5)_{ab} (\mathcal{S}_{ij} \cdot \widetilde{\mathcal{H}}) \\ &+ (\gamma^5 \gamma^\mu)_{ab} (\mathcal{S}_{ij} \cdot \widetilde{\mathcal{H}}_\mu) , \end{aligned} \quad (98)$$

This equation defines the operators $(\mathcal{A}_{ij} \cdot \mathcal{H}_\mu)$, $(\mathcal{A}_{ij} \cdot \mathcal{H}_{\mu\nu})$, $(\mathcal{S}_{ij} \cdot \mathcal{H})$, $(\mathcal{S}_{ij} \cdot \widetilde{\mathcal{H}})$, and $(\widetilde{\mathcal{H}}_\mu)$ which are appropriate for $\mathcal{N}$ -extended superspace. Finally, for all off-shell linear representations subjected to 0-brane reduction in the Coulomb gauge, and node-lowering re-definitions of fields leading to valise supermultiplets, the final form of this equation is

$$\begin{aligned} [D_{ai} , D_{bj}] &= i(\gamma^\mu)_{ab} (\mathcal{A}_{ij} \cdot \mathcal{H}_\mu) \partial_\tau + i(\gamma^\mu \wedge \gamma^\nu)_{ab} (\mathcal{A}_{ij} \cdot \mathcal{H}_{\mu\nu}) \partial_\tau \\ &+ C_{ab} (\mathcal{S}_{ij} \cdot \mathcal{H}) \partial_\tau + (\gamma^5)_{ab} (\mathcal{S}_{ij} \cdot \widetilde{\mathcal{H}}) \partial_\tau \\ &+ (\gamma^5 \gamma^\mu)_{ab} (\mathcal{S}_{ij} \cdot \widetilde{\mathcal{H}}_\mu) \partial_\tau . \end{aligned} \quad (99)$$

where the coefficients preceding the partial derivatives define the holonomy tensors acting on any valise representation of 4D, $\mathcal{N}$ -extended supersymmetry. In general these are representation depend and in general distinct for the bosonic versus fermionic fields in a supermultiplet as seen in our $\mathcal{N} = 1$ examples. Finally, it is important to note the quantities $\mathcal{H}$, $\mathcal{H}_\mu$, $\mathcal{H}_{\mu\nu}$, $\widetilde{\mathcal{H}}_\mu$, and $\widetilde{\mathcal{H}}$ in (99) are not operators. When (99) is evaluated on the bosons of a valise supermultiplet we conventionally write

$$[D_{ai}, D_{bj}] = -i2(\mathcal{B}_{aibj})\partial_\tau, \quad (100)$$

where $(\mathcal{B}_{aibj})$ denotes a set of representation dependent constants that may be decomposed in the basis in (96), when (99) is evaluated on the fermions of a valise supermultiplet we write

$$[D_{ai}, D_{bj}] = -i2(\mathcal{F}_{aibj})\partial_\tau, \quad (101)$$

where $(\mathcal{F}_{aibj})$ denotes a set of representation dependent constants that may once more be decomposed in the basis in (96).

In contrast to the objects shown in (96), the objects below

$$\begin{aligned} &(\gamma^\mu)_{ab}\delta_{ij}, \quad (\gamma^\mu \wedge \gamma^\nu)_{ab}\delta_{ij}, \\ &(\gamma^\mu)_{ab}\mathcal{S}_{ij}^{(T)}, \quad (\gamma^\mu \wedge \gamma^\nu)_{ab}\mathcal{S}_{ij}^{(T)}, \\ &C_{ab}\mathcal{A}_{ij}, \quad (\gamma^5)_{ab}\mathcal{A}_{ij}, \quad (\gamma^5\gamma^\mu)_{ab}\mathcal{A}_{ij}, \end{aligned} \quad (102)$$

are all symmetric under the exchange of the pairs of indices $ai \leftrightarrow bj$. It should be noted all of these are actually determined by the relations between the quantities in (96) and the $\text{SO}(4\mathcal{N})$ generators. In other words, a set of generators for $\text{SO}(4\mathcal{N})$ can be used to define the quantities in (102) via definitions that arise from (96).

In terms of the quantities in (102), we can write

$$\begin{aligned} \{D_{ai}, D_{bj}\} &= i(\gamma^\mu)_{ab}\delta_{ij}\mathcal{K}_\mu\partial_\tau + i(\gamma^\mu \wedge \gamma^\nu)_{ab}\delta_{ij}\mathcal{K}_{\mu\nu}\partial_\tau \\ &+ i(\gamma^\mu)_{ab}(\mathcal{S}_{ij}^{(T)} \cdot \mathcal{K}_\mu^{(T)})\partial_\tau + i(\gamma^\mu \wedge \gamma^\nu)_{ab}(\mathcal{S}_{ij}^{(T)} \cdot \mathcal{K}_{\mu\nu}^{(T)})\partial_\tau \\ &+ C_{ab}(\mathcal{A}_{ij} \cdot \mathcal{K})\partial_\tau + (\gamma^5)_{ab}(\mathcal{A}_{ij} \cdot \widetilde{\mathcal{K}})\partial_\tau \\ &+ (\gamma^5\gamma^\mu)_{ab}(\mathcal{A}_{ij} \cdot \widetilde{\mathcal{K}}_\mu)\partial_\tau, \end{aligned} \quad (103)$$

for some matrices of coefficients $\mathcal{K}_\mu$, $\mathcal{K}_{\mu\nu}$, $\mathcal{K}^{(T)}$, $\mathcal{K}_\mu^{(T)}$, $\mathcal{K}$, $\widetilde{\mathcal{K}}_\mu$, and $\widetilde{\mathcal{K}}$. From this expression it is seen the condition that the 0-brane representation should satisfy and off-shell SUSY algebra is equivalent to the requirement that only $\mathcal{K}_0$ should be non-vanishing.

The end result of the discussion in this chapter is the notion of covariant 0-brane holonomy tensors, presented in the previous chapters for the case of off-shell 4D, $\mathcal{N} = 1$ supermultiplets, can easily be extended to four dimensional theories where $\mathcal{N} > 1$.

11 Conclusion, Observations and Summary

By this work, we have been led to reason “SUSY Holography” exists between 4D, $\mathcal{N}$ -extended 0-brane SUSY valise supermultiplets on one side of a correspondence and 1D, $4\mathcal{N}$ -extended adinkra-based valise supermultiplets on the other side is due to the identity $[GL(4, \mathbb{R}) \otimes GL(\mathcal{N}, \mathbb{R})]_A = SO(4\mathcal{N})$. The 0-brane models realize the left hand side of this correspondence making manifest 4D Lorentz and $O(\mathcal{N})$ symmetries, and the adinkra models realize the right hand side making manifest connections to “Garden Algebras,” graph theory and error-correcting codes.

All our work on the adinkra substructure of 4D, $\mathcal{N} = 1$ theories inform us that all such irreducible supermultiplets appear to be in the spinor representations of $SO(4)$ which can be subject to linear restrictions that involve error-correcting codes. The spinor representations of $SO(4)$ contain an $SU(2) \otimes SU(2)$ subgroup. To distinguish between these two copies of $SU(2)$, we have in the past used the notation $SU_\alpha(2) \otimes SU_\beta(2)$ [23].

The $SU_\alpha(2)$ factor corresponds apparently to the angular momentum associated with representations in $SO(1,3)$ theories. Thus, in our work we have found the $SU_\alpha(2)$ is closely related to the spin angular momentum representations contained in the super multiplet and the corresponding adinkra network. On the other hand, the $SU_\beta(2)$ representation factor is apparently related to an extension of R-symmetry. This “extended $SU(2)$ R-symmetry” is only apparent in one dimensional valise representations and is usually broken in higher dimensional representations. This behavior is very different from what is seen in the spinning particle with extended SUSY [33].

In a classical work [33] it was shown the Hilbert space of the N -extended spinning particle leads to a spectrum of states whose spin is related to N (the number of supercharges) by the relation $spin = (1/2)N$. On the other hand, within the adinkra network approach with $N = 4$ (i.e. only four supercharges), we have found the “size” of the $SU_\alpha(2)$ representations contained in the corresponding adinkra network determines the maximum spin in the supermultiplet. In the network-based adinkra approach, changing the number of link colors in the adinkra, amounts to changing the number of space-time supercharges on the 0-brane equivalent.

This raises the very intriguing question of whether it is possible to construct a spinning particle-like action where an $O(4)$ symmetry is the one seen in the adinkra network-based approach appears?

Our next observation is the methodology we have discussed in this paper can be extended to form similar analyses of on-shell supersymmetrical theories. Owing to the fact that such theories do *not* form representations of the “Garden Algebra” as noted in [23], any such analysis is expected to more complicated than that presented with this work for the off-shell theories.

We have long been aware the definition of the L-matrices and R-matrices required by 1D, N -extended SUSY implies the quantites

$$\gamma_I = \begin{bmatrix} 0 & L_I \\ R_J & 0 \end{bmatrix} \quad (104)$$

define a set of $SO(4)$ γ matrices. Furthermore for these $SO(4)$ γ matrices, a $SO(4)$ ‘chirality’ γ^{CH} -matrix can be defined by

$$\gamma^{CH} = \begin{bmatrix} \mathbf{I}_4 & 0 \\ 0 & -\mathbf{I}_4 \end{bmatrix} \quad (105)$$

so

$$L_I = \frac{1}{2} [\mathbf{I}_8 + \gamma^{CH}] \gamma_I \quad , \quad R_I = \frac{1}{2} [\mathbf{I}_8 - \gamma^{CH}] \gamma_I \quad . \quad (106)$$

The equation in (34) implies the $\tilde{V}$ -matrices occur as projections arising from the covariant fermionic holonomy tensor $(\mathcal{F}_{ab})$. Using the equations of (4), (5), (??), and (21), the “Adinkra/ γ -matrix Equation” can be written as

$$\begin{aligned} \mathcal{P} \left[(R^\Delta)_{\hat{\Lambda} a} (L_\Delta)_b^{\hat{\Delta}} - (R^\Delta)_{\hat{\Lambda} b} (L_\Delta)_a^{\hat{\Delta}} \right] = \\ \left[(R_I)_{i^j} (L_J)_j^{\hat{k}} - (R_J)_{i^j} (L_I)_j^{\hat{k}} \right] \quad , \end{aligned} \quad (107)$$

where the projection operator simply has the effect of changing the labels on the indices according to $\mathcal{P} : (\hat{\Lambda}, \hat{\Delta}) \rightarrow (\hat{i}, \hat{k})$ and $\mathcal{P} : (a, b) \rightarrow (I, J)$.

The left hand side of (107) involves some Fierz identities of the γ -matrices for $SO(1,3)$ while on the right hand side there are the chiral parts of the $SO(4)$ generators calculated in the eight dimensional spinor representation. Identities of this type appear to be the fundamental reason SUSY Holography has worked in all our studies completed thus far. We are not aware of any prior statements in the literature on the existence of identities of this character.

The final observation we make concerns the results reported in equations (26) - (30). The matrix calculated on the 0-brane reduction side is only a 5×5 matrix as

there only exist the five representations (CS), (VS), (TS), (AVS), and (ATS)⁶. On the adinkra side, however, if one does not invoke any equivalence relations, there are 1,536 separate representations as found in [20].

A total 1,180,416 inner products would need to be calculated on the adinkra side in order to test the proposition that the only entries in (6) are either 0, 1, or $-1/3$. This is the most severe test of the concept of “SUSY Holography” proposed to date. However, if there are entries among the 1,180,416 inner products that do *not* correspond to either 0, 1, or $-1/3$, the metric on the space can be used as a filter to reject the corresponding network-based adinkras as candidates that correspond to being “shadows” of the higher dimensional supermultiplets.

Let us end by noting that the concept of supersymmetrical 0-brane holoraumy tensors does not depend on the existence of adinkra networks. The results in chapters four and five are wholly independent of the networks as all calculations in these chapter can be completed without ever knowing any properties of the networks. This means that 0-brane holoraumy tensor and the metric defined in (27) provide *new* tools for exploring the representation theory of supersymmetrical theories.

In future works, we will report on our continuing efforts, remaining conscious of a warning from Aristotle.

“The so-called Pythagoreans, who were the first to take up mathematics, not only advanced this subject, but saturated with it, they fancied that the principles of mathematics were the principles of all things.” - Aristotle

Acknowledgements

We would like to acknowledge Professors Kevin Iga, Tristan Hübsch, and Stefan Mendez-Diaz for conversations. This work was partially supported by the National Science Foundation grants PHY-0652983 and PHY-0354401. This research was also supported in part the University of Maryland Center for String & Particle Theory. Additional acknowledgment is given by M. Calkins, and D. E. A. Gates, to the Center for String and Particle Theory, as well as recognition for their participation in 2013 & 2014 SSTPRS (Student Summer Theoretical Physics Research Session) programs. Some adinkras were drawn with the aid of the *Adinkramat* © 2008 by G. Landweber. and others by T. Hübsch.

⁶The CS actually has two ‘variant representations [30], while the (VS) and (AVS) each possess a single such representation.

Appendix A: Covariant 0-Brane Bosonic Holoraumy Tensors

In this appendix, we briefly give the results for the calculations of the bosonic holoraumy tensors for four of the supermultiplets seen below,

$$\begin{aligned}
(\mathcal{B}_{ab}^{(CS)})_i^j &= C_{ab}(\gamma^{12})_i^j + i(\gamma^5)_{ab}(\gamma^{23})_i^j - i(\gamma^5\gamma^0)_{ab}(\gamma^{31})_i^j, \\
(\mathcal{B}_{ab}^{(VS)})_i^j &= -i(\gamma^5\gamma^1)_{ab}(\gamma^{23})_i^j + i(\gamma^5\gamma^2)_{ab}(\gamma^{12})_i^j - i(\gamma^5\gamma^3)_{ab}(\gamma^{31})_i^j, \\
(\mathcal{B}_{ab}^{(TS)})_i^j &= -i(\gamma^5\gamma^1)_{ab}(\gamma^{12})_i^j - i(\gamma^5\gamma^2)_{ab}(\gamma^{23})_i^j - i(\gamma^5\gamma^3)_{ab}(\gamma^{31})_i^j, \\
(\mathcal{B}_{ab}^{(TCS)})_i^j &= -i(\gamma^5\gamma^1)_{ab}(\gamma^{31})_i^j - i(\gamma^5\gamma^2)_{ab}(\gamma^{12})_i^j + i(\gamma^5\gamma^3)_{ab}(\gamma^{23})_i^j.
\end{aligned} \tag{A.1}$$

As it is not yet clear what role of the bosonic holoraumy tensors play, we have not concentrated upon them in the current work. In the future we plan to take up this question.

Appendix B:

The standard Chiral Supermultiplet formulation is given by

$$\begin{aligned}
D_a A &= \psi_a, & D_a B &= i(\gamma^5)_a^b \psi_b, \\
D_a \psi_b &= i(\gamma^\mu)_{ab}(\partial_\mu A) - (\gamma^5\gamma^\mu)_{ab}(\partial_\mu B) \\
&\quad - iC_{ab}F + (\gamma^5)_{ab}G, \\
D_a F &= (\gamma^\mu)_a^b \partial_\mu \psi_b, & D_a G &= i(\gamma^5\gamma^\mu)_a^b \partial_\mu \psi_b,
\end{aligned} \tag{B.1}$$

which leads to a Chiral Supermultiplet Valise formulation with 1D, $N = 4$ SUSY given by,

$$\begin{aligned}
D_a A &= \psi_a, & D_a B &= i(\gamma^5)_a^b \psi_b, \\
D_a F &= (\gamma^0)_a^b \psi_b, & D_a G &= i(\gamma^5\gamma^0)_a^b \psi_b, \\
D_a \psi_b &= i(\gamma^0)_{ab}(\partial_\tau A) - (\gamma^5\gamma^0)_{ab}(\partial_\tau B) \\
&\quad - iC_{ab}(\partial_\tau F) + (\gamma^5)_{ab}(\partial_\tau G).
\end{aligned} \tag{B.2}$$

Next in a similar manner we begin with the 4D, $N = 1$ Vector Supermultiplet formulation

$$\begin{aligned}
D_a A_\mu &= (\gamma_\mu)_a^b \lambda_b, \\
D_a \lambda_b &= -i\frac{1}{2}(\gamma^\mu\gamma^\nu)_{ab}F_{\mu\nu} + (\gamma^5)_{ab}d, \\
D_a d &= i(\gamma^5\gamma^\mu)_a^b \partial_\mu \lambda_b,
\end{aligned} \tag{B.3}$$

which leads to a Vector Supermultiplet Valise formulation with 1D, $N = 4$ SUSY given by,

$$\begin{aligned} D_a A_m &= (\gamma_m)_a{}^b \lambda_b \quad , \quad D_a d = i(\gamma^5 \gamma^0)_a{}^b \lambda_b \quad , \\ D_a \lambda_b &= -i(\gamma^0 \gamma^m)_{ab} (\partial_\tau A_m) + (\gamma^5)_{ab} (\partial_\tau d) \quad . \end{aligned} \quad (B.4)$$

For the 4D, $N = 1$ Tensor Supermultiplet formulation

$$\begin{aligned} D_a \varphi &= \chi_a \quad , \\ D_a B_{\mu\nu} &= -\frac{1}{4}([\gamma_\mu, \gamma_\nu])_a{}^b \chi_b \quad , \\ D_a \chi_b &= i(\gamma^\mu)_{ab} \partial_\mu \varphi - (\gamma^5 \gamma^\mu)_{ab} \epsilon_\mu{}^{\rho\sigma\tau} \partial_\rho B_{\sigma\tau} \quad . \end{aligned} \quad (B.5)$$

leads to a Tensor Supermultiplet Valise formulation with 1D, $N = 4$ SUSY given by,

$$\begin{aligned} D_a \varphi &= \chi_a \quad , \quad D_a B_{mn} = -\frac{1}{4}([\gamma_m, \gamma_n])_a{}^b \chi_b \quad , \\ D_a \chi_b &= i(\gamma^0)_{ab} \partial_\tau \varphi - i\frac{1}{2}(\gamma^0 [\gamma^m, \gamma^n])_{ab} \partial_\tau B_{mn} \quad . \end{aligned} \quad (B.6)$$

Finally, there exist a second supermultiplet which contains a spin-1 gauge field. The component fields shown in (B.1) and (B.2) are appropriate since the gauge field is a normal vector. However, it is also possible to introduce a axial Vector Supermultiplet (AVS) with component fields U_μ , $\tilde{\lambda}_a$, and $\tilde{d}$. The main difference with the VS is here the component field U_μ corresponds to an axial vector. Accordingly, the D-equations are changed to the forms

$$\begin{aligned} D_a U_\mu &= i(\gamma^5 \gamma_\mu)_a{}^b \tilde{\lambda}_b \quad , \\ D_a \tilde{\lambda}_b &= \frac{1}{2}(\gamma^5 \gamma^\mu \gamma^\nu)_{ab} F_{\mu\nu}(U) + i C_{ab} \tilde{d} \quad , \\ D_a \tilde{d} &= -(\gamma^\mu)_a{}^b \partial_\mu \tilde{\lambda}_b \quad , \end{aligned} \quad (B.7)$$

where $F_{\mu\nu}(U)$ simply indicates the ‘curl’ of the U -gauge field. So an axial Vector Supermultiplet Valise formulation with 1D, $N = 4$ SUSY is given by,

$$\begin{aligned} D_a U_m &= i(\gamma^5 \gamma_m)_a{}^b \tilde{\lambda}_b \quad , \quad D_a \tilde{d} = -(\gamma^0)_a{}^b \tilde{\lambda}_b \quad , \\ D_a \tilde{\lambda}_b &= (\gamma^5 \gamma^0 \gamma^m)_{ab} (\partial_\tau U_m) + i C_{ab} (\partial_\tau \tilde{d}) \quad . \end{aligned} \quad (B.8)$$

On comparing (B.3) with (B.7), it is seen the latter corresponds applying a set of replacements in the former according to $V \rightarrow U$, $\lambda_a \rightarrow -i(\gamma^5)_a{}^b \tilde{\lambda}_b$, and $d \rightarrow \tilde{d}$. In turn, this observation implies the possibility of constructing an axial Tensor Supermultiplet (ATS) via replacements in the (B.6) according to: $B \rightarrow \tilde{B}$, $\chi_a \rightarrow -i(\gamma^5)_a{}^b \tilde{\chi}_b$, and $\varphi \rightarrow \tilde{\varphi}$.

Let us emphasize the bosonic and fermionic functions appearing in (B.2), (B.4), (B.6), and (B.7) depend solely on a single real continuous parameter τ . Though

these field variable have indices, these should be regarded as isospin indices as there are no spatial coordinate in these four equations. In a similar manner, the matrices multiplying such indices are also to be thought of as linear operators acting in isospin space. It should be noted it is consistent to assign to all the bosonic fields in (B.2), (B.4), (B.6), and (B.7) the same engineering dimensions. Likewise this can also be done for all the fermionic fields in the same four equations, but of course the engineering dimension for the bosonic fields are distinct from that of the fermionic fields.

Appendix C: Detailed Derivation

In this appendix, we provide a detailed derivation of the steps led from the equations in (36) to those in (37). As indicated in (36) we begin with the equations

$$\begin{aligned} D_a A_\mu &= (\gamma_\mu)_a{}^b \lambda_b \quad , \\ D_a \lambda_b &= -i \frac{1}{4} ([\gamma^\mu, \gamma^\nu])_{ab} (\partial_\mu A_\nu - \partial_\nu A_\mu) + (\gamma^5)_{ab} d \quad , \\ D_a d &= i (\gamma^5 \gamma^\mu)_a{}^b \partial_\mu \lambda_b \quad , \end{aligned} \tag{C.1}$$

as our starting point. The first of these sets of equations in (C.1) simply gives us

$$\begin{aligned} D_a A_0 &= (\gamma_0)_a{}^b \lambda_b \quad , \quad D_a A_1 = (\gamma_1)_a{}^b \lambda_b \quad , \\ D_a A_2 &= (\gamma_2)_a{}^b \lambda_b \quad , \quad D_a A_3 = (\gamma_3)_a{}^b \lambda_b \quad , \end{aligned} \tag{C.2}$$

and the second set yields

$$\begin{aligned} D_a \lambda_b &= -i \frac{1}{2} ([\gamma^0, \gamma^1])_{ab} (\partial_0 A_1 - \partial_1 A_0) - i \frac{1}{2} ([\gamma^1, \gamma^2])_{ab} (\partial_1 A_2 - \partial_2 A_1) \\ &\quad - i \frac{1}{2} ([\gamma^0, \gamma^2])_{ab} (\partial_0 A_2 - \partial_2 A_0) - i \frac{1}{2} ([\gamma^2, \gamma^3])_{ab} (\partial_2 A_3 - \partial_3 A_2) \\ &\quad - i \frac{1}{2} ([\gamma^0, \gamma^3])_{ab} (\partial_0 A_3 - \partial_3 A_0) - i \frac{1}{2} ([\gamma^3, \gamma^1])_{ab} (\partial_3 A_1 - \partial_1 A_3) \\ &\quad + (\gamma^5)_{ab} d \quad . \end{aligned} \tag{C.3}$$

Next in (C.1) we delete entirely the first equation and go to the Coulomb gauge in (C.2). This leaves us with

$$D_a A_1 = (\gamma_1)_a{}^b \lambda_b \quad , \quad D_a A_2 = (\gamma_2)_a{}^b \lambda_b \quad , \quad D_a A_3 = (\gamma_3)_a{}^b \lambda_b \quad , \tag{C.4}$$

and in (C.3) we use both the gauge condition and the restrictions on the coordinate dependence to find

$$\begin{aligned} D_a \lambda_b &= +i \frac{1}{2} ([\gamma^2, \gamma^0])_{ab} (\partial_0 A_2) + i \frac{1}{2} ([\gamma^2, \gamma^1])_{ab} (\partial_1 A_2) \\ &\quad + i \frac{1}{2} ([\gamma^3, \gamma^0])_{ab} (\partial_0 A_3) + i \frac{1}{2} ([\gamma^3, \gamma^1])_{ab} (\partial_1 A_3) \\ &\quad + (\gamma^5)_{ab} d - i \frac{1}{2} ([\gamma^0, \gamma^1])_{ab} (\partial_0 A_1) \quad . \end{aligned} \tag{C.5}$$

With a bit more algebra

$$\begin{aligned}
D_a \lambda_b = & + i(\gamma^2 \gamma^0)_{ab} (\partial_0 A_2) + i(\gamma^2 \gamma^1)_{ab} (\partial_1 A_2) \\
& + i(\gamma^3 \gamma^0)_{ab} (\partial_0 A_3) + i(\gamma^3 \gamma^1)_{ab} (\partial_1 A_3) \\
& + (\gamma^5)_{ab} d - i(\gamma^0 \gamma^1)_{ab} \widehat{F} \ ,
\end{aligned} \tag{C.6}$$

where we have introduced the notation

$$\widehat{F} = (\partial_0 A_1) \ . \tag{C.7}$$

Finally the last equations of (C.1) become

$$D_a d = i(\gamma^5 \gamma^0)_a{}^b \partial_0 \lambda_b + i(\gamma^5 \gamma^1)_a{}^b \partial_1 \lambda_b \ . \tag{C.8}$$

The latter two equations of (C.4), along with (C.6), and (C.7) are covariant with respect to the two dimensional Lorentz symmetry of the space with coordinates x^0 and x^1 . For (C.6) and (C.7) we must take into consideration the Coulomb gauge condition.

We can make one final change of notation

$$A_2 \equiv \widehat{A} \ , \ A_3 \equiv \widehat{B} \ , \ \lambda_a \equiv \widehat{\psi}_a \ , \ d \equiv \widehat{G} \ , \tag{C.9}$$

to obtain

$$\begin{aligned}
D_a \widehat{A} &= (\gamma_2)_a{}^b \widehat{\psi}_b \ , \ D_a \widehat{B} = (\gamma_3)_a{}^b \widehat{\psi}_b \ , \\
D_a \widehat{\psi}_b &= i(\gamma^2 \gamma^0)_{ab} (\partial_0 \widehat{A}) + i(\gamma^2 \gamma^1)_{ab} (\partial_1 \widehat{A}) \\
&\quad + i(\gamma^3 \gamma^0)_{ab} (\partial_0 \widehat{B}) + i(\gamma^3 \gamma^1)_{ab} (\partial_1 \widehat{B}) \\
&\quad - i(\gamma^0 \gamma^1)_{ab} \widehat{F} + (\gamma^5)_{ab} \widehat{G} \ , \\
D_a \widehat{F} &= (\gamma_1)_a{}^b (\partial_0 \widehat{\psi}_b) - (\gamma_0)_a{}^b (\partial_1 \widehat{\psi}_b) \ , \\
D_a \widehat{G} &= i(\gamma^5 \gamma^0)_a{}^b (\partial_0 \widehat{\psi}_b) + i(\gamma^5 \gamma^1)_a{}^b (\partial_1 \widehat{\psi}_b) \ .
\end{aligned} \tag{C.10}$$

This is a formulation of the $\mathcal{N} = 2$ twisted chiral supermultiplet (TCS) in two dimensions. The process above is precisely the one that led to the discovery of the twisted chiral supermultiplet.

Appendix D: Comparison of Defining Conditions

The equations on the first line of (39) can be used to derive a first order spinorial differential equation on A and B . We note

$$D_a B = i(\gamma^5)_a{}^b \psi_b \Rightarrow \psi_b = -i(\gamma^5)_b{}^c D_c B \ , \tag{D.1}$$

and this final result can be substituted into the first equation in (40) to derive

$$D_a A + i(\gamma^5)_a{}^b D_b B = 0 \quad . \quad (D.2)$$

By using similar arguments based on (37) we find a first order spinorial differential equation imposed upon $\widehat{A}$, and $\widehat{B}$

$$D_a \widehat{B} = (\gamma_3)_a{}^b \widehat{\psi}_b \Rightarrow \widehat{\psi}_b = (\gamma_3)_b{}^c D_c \widehat{B} \quad , \quad (D.3)$$

which leads to

$$D_a \widehat{A} - (\gamma_2 \gamma_3)_a{}^b D_b \widehat{B} = 0 \quad . \quad (D.4)$$

The condition in (D.2) corresponds to the real formulation of what is known as the “chirality condition” and in a similar manner (D.4) corresponds to the real formulation of what is known as the “twisted chirality condition.”

The equation in (D.1) implies

$$\begin{aligned} \rightarrow D_b(A + iB) &= [(I)_b{}^c - (\gamma^5)_b{}^c] \psi_c \quad , \\ \rightarrow \frac{1}{2} [(I) + (\gamma^5)]_a{}^b D_b(A + iB) &= 0 \quad , \\ \rightarrow \frac{1}{2} \begin{bmatrix} D_1 + iD_4 \\ D_2 - iD_3 \\ i(D_2 - iD_3) \\ -i(D_1 + iD_4) \end{bmatrix} (A + iB) &= 0 \quad , \end{aligned} \quad (D.5)$$

so there are only two linearly independent defining conditions

$$\left[D_1 + iD_4 \right] (A + iB) = 0 \quad , \quad \left[D_2 - iD_3 \right] (A + iB) = 0 \quad , \quad (D.6)$$

and of course the complex conjugates of these equations.

The equation in (D.4) implies

$$\begin{aligned} D_a \widehat{A} &= \widehat{\psi}_a \quad , \quad D_a \widehat{B} = -(\gamma_2 \gamma_3)_a{}^b \widehat{\psi}_b \quad , \\ \rightarrow D_b(\widehat{A} + i\widehat{B}) &= [(I)_b{}^c - i(\gamma^2 \gamma^3)_b{}^c] \widehat{\psi}_b \quad , \\ \rightarrow \frac{1}{2} [(I) + i(\gamma^2 \gamma^3)]_a{}^b D_b(\widehat{A} + i\widehat{B}) &= 0 \quad , \\ \rightarrow \frac{1}{2} \begin{bmatrix} D_1 + iD_4 \\ D_2 + iD_3 \\ -i(D_2 + iD_3) \\ -i(D_1 + iD_4) \end{bmatrix} (\widehat{A} + i\widehat{B}) &= 0 \quad . \end{aligned} \quad (D.7)$$

Once more there are only two linearly independent defining conditions,

$$\left[D_1 + iD_4 \right] (\widehat{A} + i\widehat{B}) = 0 \quad , \quad \left[D_2 + iD_3 \right] (\widehat{A} + i\widehat{B}) = 0 \quad , \quad (D.8)$$

and of course the complex conjugates of these equations also. The only difference between (D.6) and (D.8) is the replacement $D_3 \rightarrow -D_3$. If we write the two equations in (D.6) in the forms of

$$\bar{D}_+(A + iB) = 0 \quad , \quad \bar{D}_-(A + iB) = 0 \quad , \quad (D.9)$$

utilizing the 2D complex light supercovariant derivatives $\bar{D}_+$, and $\bar{D}_-$ then we find (D.8) takes the form

$$\bar{D}_+(\hat{A} + i\hat{B}) = 0 \quad , \quad D_-(\hat{A} + i\hat{B}) = 0 \quad , \quad (D.10)$$

which is in accord with the definitions of the chiral and twisted chiral 2D, $\mathcal{N} = 2$ supermultiplets given in 1984.

Appendix E:

In the work of [23], we introduced shorthand notations for 4×4 matrices $\alpha^1, \alpha^2, \alpha^3, \beta^1, \beta^2, \beta^3$ where

$$\begin{aligned} \alpha^1 &= \sigma^2 \otimes \sigma^1 & , & & \alpha^2 &= \mathbf{I}_2 \otimes \sigma^2 & , & & \alpha^3 &= \sigma^2 \otimes \sigma^3 & , \\ &= i(3)_b(14)(23) & , & & &= i(5)_b(12)(34) & , & & &= i(9)_b(13)(24) & , \\ \beta^1 &= \sigma^1 \otimes \sigma^2 & , & & \beta^2 &= \sigma^2 \otimes \mathbf{I}_2 & , & & \beta^3 &= \sigma^3 \otimes \sigma^2 & , \\ &= i(5)_b(14)(23) & , & & &= i(3)_b(13)(24) & , & & &= i(9)_b(12)(34) & . \end{aligned} \quad (E.1)$$

where the outer product conventions are in [23] and the Boolean factor/permutation conventions are in [21]. These matrices satisfy the identities

$$\begin{aligned} \alpha^{\hat{I}} \alpha^{\hat{K}} &= \delta^{\hat{I}\hat{K}} \mathbf{I}_4 + i \epsilon^{\hat{I}\hat{K}\hat{L}} \alpha^{\hat{L}} & , & & \beta^{\hat{I}} \beta^{\hat{K}} &= \delta^{\hat{I}\hat{K}} \mathbf{I}_4 + i \epsilon^{\hat{I}\hat{K}\hat{L}} \beta^{\hat{L}} & , \\ \text{Tr}(\alpha^{\hat{I}}) &= \text{Tr}(\beta^{\hat{I}}) = 4 & , & & [\alpha^{\hat{I}} , \beta^{\hat{K}}] &= 0 & . \end{aligned} \quad (E.2)$$

These will be useful in the following discussion of this chapter.

Now we can use the adinkra in (Adnk-A) to find the associated L-matrices (and R-matrices which are simply the transposed versions of the L-matrices). Using the notation in (E.1), and the rainbow assignment of (black, green, red, blue) colors to corresponding link numbers (1, 2, 3, 4) respectively, for the L-matrices yields

$$\begin{aligned} L_1^{(Adnk-A)} &= \mathbf{I}_4 & , & & L_2^{(Adnk-A)} &= i\beta^3 & , \\ L_3^{(Adnk-A)} &= -i\beta^2 & , & & L_4^{(Adnk-A)} &= -i\beta^1 & , \end{aligned} \quad (E.3)$$

associated with (Adnk-A). These imply the corresponding $\tilde{V}$ matrices associated with (Adnk-A) take the forms

$$\begin{aligned}\tilde{V}_{12}^{(\text{Adnk-A})} &= i\beta^3, & \tilde{V}_{23}^{(\text{Adnk-A})} &= i\beta^1, & \tilde{V}_{34}^{(\text{Adnk-A})} &= i\beta^3, \\ \tilde{V}_{13}^{(\text{Adnk-A})} &= -i\beta^2, & \tilde{V}_{24}^{(\text{Adnk-A})} &= -i\beta^2, \\ \tilde{V}_{14}^{(\text{Adnk-A})} &= -i\beta^1,\end{aligned}\tag{E.4}$$

and these equations further imply

$$\begin{aligned}\tilde{V}_{12}^{(\text{Adnk-A})} &= \tilde{V}_{34}^{(\text{Adnk-A})}, & \tilde{V}_{23}^{(\text{Adnk-A})} &= -\tilde{V}_{14}^{(\text{Adnk-A})}, \\ \tilde{V}_{13}^{(\text{Adnk-A})} &= \tilde{V}_{24}^{(\text{Adnk-A})}.\end{aligned}\tag{E.5}$$

By use of the adinkra in (Adnk-B), it is seen one only needs to make the replacement of $L_3 \rightarrow -L_3$ to find

$$\begin{aligned}L_1^{(\text{Adnk-B})} &= \mathbf{I}_4, & L_2^{(\text{Adnk-B})} &= i\beta^3, \\ L_3^{(\text{Adnk-B})} &= i\beta^2, & L_4^{(\text{Adnk-B})} &= -i\beta^1,\end{aligned}\tag{E.6}$$

and this leads to the corresponding $\tilde{V}$ matrices given by

$$\begin{aligned}\tilde{V}_{12}^{(\text{Adnk-B})} &= i\beta^3, & \tilde{V}_{23}^{(\text{Adnk-B})} &= -i\beta^1, & \tilde{V}_{34}^{(\text{Adnk-B})} &= -i\beta^3, \\ \tilde{V}_{13}^{(\text{Adnk-B})} &= i\beta^2, & \tilde{V}_{24}^{(\text{Adnk-B})} &= -i\beta^2, \\ \tilde{V}_{14}^{(\text{Adnk-B})} &= -i\beta^1,\end{aligned}\tag{E.7}$$

and these equations further imply

$$\begin{aligned}\tilde{V}_{12}^{(\text{Adnk-B})} &= -\tilde{V}_{34}^{(\text{Adnk-B})}, & \tilde{V}_{23}^{(\text{Adnk-B})} &= \tilde{V}_{14}^{(\text{Adnk-B})}, \\ \tilde{V}_{13}^{(\text{Adnk-B})} &= -\tilde{V}_{24}^{(\text{Adnk-B})}.\end{aligned}\tag{E.8}$$

We can further see

$$\begin{aligned}\tilde{V}_{12}^{(\text{Adnk-A})} &= -\tilde{V}_{12}^{(\text{Adnk-B})}, & \tilde{V}_{13}^{(\text{Adnk-A})} &= -\tilde{V}_{13}^{(\text{Adnk-B})}, \\ \tilde{V}_{14}^{(\text{Adnk-A})} &= \tilde{V}_{14}^{(\text{Adnk-B})}.\end{aligned}\tag{E.9}$$

From the results shown in (E.8) and (E.9) it is clear

$$\tilde{V}_{IJ}^{(\text{Adnk-A})} \tilde{V}_{KL}^{(\text{Adnk-B})} - \tilde{V}_{KL}^{(\text{Adnk-B})} \tilde{V}_{IJ}^{(\text{Adnk-A})} \neq 0.\tag{E.10}$$

References

- [1] M. Faux, S. J. Gates Jr., “Adinkras: A Graphical Technology for Supersymmetric Representation Theory,” *Phys. Rev.* **D71** (2005) 065002, [hep-th/0408004v1].
- [2] C. F. Doran, M. G. Faux, S. J. Gates, Jr., T. Hübsch, K. M. Iga, and G. D. Landweber, “Relating Doubly-Even Error-Correcting Codes, Graphs, and Irreducible Representations of N -Extended Supersymmetry,” in “Discrete and Computational Mathematics,” p. 53–71, eds. F. Liu et al., (Nova Science Pub., Inc., Hauppauge, 2008), [arXiv:0806.0051].
- [3] C. F. Doran, M. G. Faux, S. J. Gates, Jr., T. Hübsch, K. M. Iga, G. D. Landweber, and R. L. Miller, “Codes and Supersymmetry in One Dimension,” *Adv. in Theor. and Math. Phys.* **15** (2011) 1909-1970, [arXiv:1108.4124].
- [4] C. F. Doran, M. G. Faux, S. J. Gates, Jr., T. Hübsch, K. M. Iga, G. D. Landweber, and R. L. Miller, “Topology Types of Adinkras and the Corresponding Representations of N -Extended Supersymmetry,” Univ. of Maryland preprint UMDEPP-08-010, State University of New York - Oneonta SUNY-O-667 e-Print: arXiv:0806.0050, unpublished.
- [5] C. F. Doran, M. G. Faux, S. J. Gates, Jr., T. Hübsch, K. M. Iga, and G. D. Landweber, “On graph-theoretic identifications of Adinkras, supersymmetry representations and superfields,” *Int. J. Mod. Phys.* **A22** (2007) 869-930, [math-ph/0512016].
- [6] Z. Kuznetsova, M. Rojas and F. Toppan, “Classification of irreps and invariants of the N -extended supersymmetric quantum mechanics,” *JHEP* **03** (2006) 098, [hep-th/0511274].
- [7] Z. Kuznetsova, and F. Toppan, “Refining the classification of the irreps of the 1D N -Extended Supersymmetry,” *Mod. Phys. Lett.* **A23** (2008) 37-51, e-Print: hep-th/0701225.
- [8] Z. Kuznetsova, and F. Toppan, “Decomposition and oxidation of the N -extended supersymmetric quantum mechanics multiplets,” *Int. J. Mod. Phys.* **A23** (2008) 3947-3962 e-Print: arXiv:0712.3176 [hep-th].
- [9] Z. Kuznetsova, and F. Toppan, “D-module Representations of $N=2,4,8$ Superconformal Algebras and Their Superconformal Mechanics,” *J. Math. Phys.* **53** (2012) 043513, e-Print: arXiv:1112.0995 [hep-th].

- [10] M. Gonzales, K. Iga, S. Khodaei, and F. Toppan, “Pure and entangled $N=4$ linear supermultiplets and their one-dimensional sigma-models,” *J. Math. Phys.* **53** (2012) 103513, e-Print: arXiv:1204.5506 [hep-th]. .
- [11] S. J. Gates Jr., and L. Rana, “On Extended Supersymmetric Quantum Mechanics,” UMDEPP 93-194 (1994), unpublished.
- [12] S. J. Gates Jr., and L. Rana, “Ultra-Multiplets: A New Representation of Rigid 2D, $N = 8$ Supersymmetry,” *Phys. Lett.* **B342** (1995) 132-137.
- [13] S. J. Gates Jr., and L. Rana, “A Theory of Spinning Particles for Large N -extended Supersymmetry (I),” *Phys. Lett.* **B352** (1995) 50, arXiv [hep-th:9504025].
- [14] S. J. Gates Jr., and L. Rana, “A Theory of Spinning Particles for Large N -extended Supersymmetry (II),” *ibid. Phys. Lett.* **B369** (1996) 262, arXiv [hep-th:9510151].
- [15] S. J. Gates Jr., and L. Rana, “Tuning the RADIO to the off-shell 2-D Fayet hypermultiplet problem,” arXiv [hep-th:9602072], unpublished.
- [16] S. J. Gates, Jr., W. D. Linch, III, J. Phillips and L. Rana, *Grav. Cosmol.* **8** (2002) 96, arXiv [hep-th/0109109].
- [17] S. J. Gates, Jr., W. D. Linch, III, J. Phillips, “When Superspace Is Not Enough,” Univ. of Md Preprint # UMDEPP-02-054, Caltech Preprint # CALT-68-2387, arXiv [hep-th:0211034], unpublished.
- [18] S. J. Gates Jr., T. Hübsch, and K. Stiffler, “On Clifford-Algebraic ”Holonomy”, Dimensional Extension, and SUSY Holography,” Univ. of MD Preprint UMDEPP-014-007, arXiv:1409.4445 [hep-th], unpublished.
- [19] R. Jackiw, and C. Rebbi, “Spin from Isospin in a Gauge Theory” *Phys. Rev. Lett.* **36** (1976) 1116.
- [20] I. Chappell, II, S. J. Gates, Jr, and T. Hübsch, “Adinkra (In)Equivalence From Coxeter Group Representations: A Case Study,” *Int. J. Mod. Phys.* **A29** (2014) 1450029, e-Print: arXiv:1210.0478 [hep-th].
- [21] S. J. Gates, Jr., S. Randall, and K. Stiffler, “Reduction Redux of Adinkras,” *Int.J.Mod.Phys.* **A29** (2014) 13, 1450070, e-Print: arXiv:1312.2000 [hep-th], DOI: 10.1142/S0217751X14500705.

- [22] C. Doran, K. Iga, G. Landweber, and S. Mendez-Diez “Geometrization of N-Extended 1-Dimensional Supersymmetry Algebras,” (Alberta U., Dept. Math. Stat. Sci.). Nov 15, 2013. 69 pp. e-Print: arXiv:1311.3736 [hep-th].
- [23] S.J. Gates, Jr. J. Gonzales, B. MacGregor, J. Parker, R. Polo-Sherk, V.G.J. Rodgers and L. Wassink, “4D, $N = 1$ Supersymmetry Genomics (I),” JHEP **0912**, 008 (2009), e-Print: arXiv:0902.3830 [hep-th].
- [24] S. J. Gates, Jr., “The Search for Elementarity Among Off-Shell SUSY Representations,” The Korean Institute for Advanced Studies (KIAS) Newsletter, Vol. **5** 2012, ppg 19 - 27, Seoul, Korea.
- [25] S. J. Gates, Jr., T. Hübsch, and K. Stiffler, “Adinkras and SUSY Holography: Some explicit examples,” Int. J. Mod. Phys. **A29** (2014) 1450041, e-Print: arXiv:1208.5999 [hep-th].
- [26] S. J. Gates, Jr., “Superspace Formulation of New Nonlinear Sigma Models,” Nucl. Phys. **B238** (1984) 349.
- [27] S. J. Gates, Jr., C. M. Hull, M. Rőcek, “Twisted Multiplets and New Supersymmetric Nonlinear Sigma Models,” Nucl. Phys. **B248** (1984) 157.
- [28] M. Calkins, D. E. A. Gates, S. J. Gates, B. McPeak, “Is it possible to embed a 4D, $N=4$ supersymmetric vector multiplet within a completely off-shell adinkra hologram?,” JHEP **1405** (2014) 057, e-Print: arXiv:1402.5765 [hep-th], DOI: 10.1007/JHEP05(2014)057.
- [29] J. Wess, Lectures given at the Bonn Summer School 1974; P. Fayet, “Fermi-Bose Hypersymmetry,” Nucl. Phys. **B113** (1976) 135.
- [30] S. J. Gates, Jr., and W. Siegel, Nucl. Phys. **B187** (1981) 389.
- [31] S. J. Gates Jr., “*Super P-form Gauge Superfields*,” Nucl. Phys. **B184** 381 (1981).
- [32] S. J. Gates, Jr., ”Symbols of Power,” *Physics World*, Vol. 23, No 6, June 2010, pp. 34 - 39.
- [33] P. S. Howe, S. Penati, M. Pernici, and P. K. Townsend, “Wave Equations for Arbitrary Spin From Quantization of the Extended Supersymmetric Spinning Particle,” Phys. Lett. **B215** (1988) 555.